Secondary prevention is:

- a) creating healthy environmental conditions, controlling risk factors
- b) using diagnostic and clinic methods to prevent the disease, screening
- c) is applied during therapeutic procedures
- d) is applied after disease so as to prevent further attacks

In the lung cancer are important heavy metals:

- a) no heavy metals are carcinogenic
- b) only mercury is the factor supporting lung cancer
- c) iron and its substances
- d) cadmium, chromium, nickel, arsenic

Pattern how to account BMI:

- a) weight/height
- b) weight²/height
- c) weight/height²
- d) height (minus) weight

Obesity of the I. first grade is:

- a) BMI 20-25
- b) BMI 25-30
- c) BMI less then 20
- d) BMI 30-35

Energetic value of 1 g of proteins is:

- a) 17 KJoules
- b) 39 KJoules
- c) 17 kcal
- d) 39 kcal

Retrospective cohort studies you can use only in limited contents where you know exposure and health effect from past history:

- a) biochemical studies
- b) occupational studies and occupational cancer studies
- c) acute respiratory morbidity study
- d) cross-sectional studies

Sideropenic anaemia is one of the symptoms of:

- a) low magnesium and calcium intake
- b) low B₁₂ vitamin intake
- c) low iron intake
- d) low fluoride intake

Hypertension as a risk of cardiovascular disease is factor:

- a) internal, modifiable, third line of importance
- b) external, modifiable, first line of importance
- c) internal modifiable, fist line of importance
- d) internal, non-modifiable, first line of importance

.Find **correct** answer:

- a) ultrasound is the sound with frequency higher than 20 000 KiloHertz
- b) tryptophane belongs to the group of "limited amino acids"
- c) period of child age from 1-3 years old is called preschool age
- d) vaccination is secondary prevention

.Find **correct** answer:

- a) salt iodination is secondary prevention
- b) PAP smears in gynaecology is tertiary prevention
- c) asbestos is not carcinogenic
- d) leukaemia belongs to the stochastic effect