

Railway stations vs. thermophilous species (example from Eastern Slovakia)

JANA MÁJEKOVÁ¹, VLADIMÍR JEHLÍK² & MARICA ZALIBEROVÁ¹

¹Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovak Republic; jana.majekova@savba.sk, maria.zaliberova@savba.sk

²V Lesíčku 1, 150 00 Praha 5 – Smíchov, Czech Republic

Májeková J., Jehlík V. & Zaliberová M. (2016): Railway stations vs. thermophilous species (example from Eastern Slovakia) – Thaiszia – J. Bot. 26 (2): 173-188. – ISSN 1210-0420.

Abstract: Railway habitats host a multitude of plant taxa – both native and alien, including threatened, synanthropic and invasive species. Thermophilous species are also often present at many railway stations due to the high temperature of rails and gravel. The aims of this study are: (i) to document past and current vascular plant flora identified at Košice and Spišská Nová Ves railway stations in Eastern Slovakia (ii) to compare species composition at these stations and (iii) to evaluate species' temperature preferences at the different climatic regions (at Košice, Spišská Nová Ves, Plaveč and Čierna nad Tisou stations). Recorded plant taxa were; 207 at Košice railway station and 204 at Spišská Nová Ves. Archaeophytes prevailed over neophytes at both stations and the most frequent plant families were *Asteraceae*, *Poaceae* and *Brassicaceae*. Despite the different climates, similar species structure was identified at all studied railway stations.

Keywords: vascular plants, distribution, climate, species thermophility, Ellenberg indicator values, Borhidi indicator values, railway, Slovakia.

Introduction

Anthropogenic influences have a significant role in species spread throughout a country. In addition to deliberate plant and animal introduction for agricultural, medical and ornamental purposes, there is also accidental spread by increasing

globalization, international traffic and trade (VAN KLEUNEN et al. 2015). Plant diasporas are imported on road vehicles and with transported material, so railway has become a major factor in flora synanthropization (RABITSCH & ESSL 2006, WIŁKOMIRSKI et al. 2012). JEHLÍK (1998, 2013) pioneered research in plant spread by railway and ship traffic in Slovakia and in former Czechoslovakia. His work revealed that freight transport was more important than passenger traffic in spreading adventives.

Apart from Jejhík, only marginal attention has been devoted to flora and vegetation research at Slovak railway habitats (e.g. SCHIDLAY 1944, FRANTOVA 1947, ELIÁŠ 1977, 1979a, b, c, 1981, JEHLÍK & DOSTÁLEK 1989, 2008, ELIÁŠ jun. 2011, ELIÁŠ jun. et al. 2013, JEHLÍK et al. 2013, MÁJEKOVÁ et al. 2013, 2014, ZALIBEROVÁ & MÁJEKOVÁ 2014 and MÁJEKOVÁ & LIMÁNEK 2016).

Study on this topic in adjacent Central European countries has also been performed in the Czech Republic (JEHLÍK 1986, 1995, HOSKOVEC 2009, RŮŽIČKA & KOBLÍŽEK 2009, STAŘECKÁ 2010), Poland (PISKORZ & CZARNA 2006, NOWINSKA & CZARNA 2008, GALERA et al. 2012, 2014), Austria (HOHLA et al. 2000, 2002, 2005), Ukraine (SHEVERA et al. 2015) and Germany (cf. www.ruderal-vegetation.de epub/#eisenbahn). BRANDES (1983) concentrated on flora and vegetation at Central European railway stations and published his Bibliography on railway vegetation in Europe (BRANDES 2008).

Although railway habitats are often neglected by botanists, they are very interesting biotopes. JEHLÍK (1986, 1998) characterised synanthropic stands in the following adjacent railway areas: (1) many fields without vegetation – with reduced native species competition; (2) very young soils; (3) drying stands; (4) temperature elevated by dark substrate surface and (5) constant anthropogenic impacts on the soil surface.

Railyards, platforms and banks offer a refugium for many species; especially for both native and alien synanthropic plants, but threatened species also find these conditions suitable (cf. HUBER 1992, HOHLA et al. 2000, 2002, 2005, NOWIŃSKA & CZARNA 2008, MÁJEKOVÁ et al. 2014, MÁJEKOVÁ & LIMÁNEK 2016). These sites also provide a convenient corridor for alien and invasive plant spread (JEHLÍK 1995, RABITSCH & ESSL 2006), where some species bind to these biotopes in their initial introduction to a new country (cf. ELIÁŠ jun. 2011, ZALIBEROVÁ & MÁJEKOVÁ 2014). Thermophilous species are especially suited to the extreme train rail and gravel temperatures, because soil temperature increase above air temperature enhances their germination (BRANDES 1983); and this is also noted in reports of several thermophilous plants at railway stations in different countries (ELIÁŠ 1977, BRANDES 1983, ELIÁŠ jun. et al. 2013, GALERA et al. 2014, MÁJEKOVÁ et al. 2014, MÁJEKOVÁ & LIMÁNEK 2016).

The above research inspired us to investigate the extent to which the proportional representation of thermophilous species is dependent on railway station climate. Here, we chose the following 4 railway stations in Eastern Slovakia in different climatic regions, altitudes and phytogeographical districts: Košice, Spišská Nová Ves, Plaveč and Čierna nad Tisou. In addition to Jejhík's past railway biotope research in Eastern Slovakia (JEHLÍK & DOSTÁLEK 2008, JEHLÍK et

al. in press), he also collected data from Košice and Spišská Nová Ves stations, but his species lists are incomplete. Our comprehensive field-work therefore detailed the flora at these two stations to complete the study. In addition, Plaveč station species composition was published by MÁJEKOVÁ & LIMÁNEK (2016), and our study of Čierna nad Tisou populations is in press (JEHLIK et al. in press).

The aims of this study are: (1) to document and evaluate past and present vascular plant flora at Košice and Spišská Nová Ves railway stations; (2) to compare species composition at these sites and (3) to compare species' temperature preferences at Košice, Spišská Nová Ves, Plaveč and Čierna nad Tisou Eastern Slovak railway stations.

Material and methods

Study area

Field research was performed at Košice and Spišská Nová Ves railway stations. These are situated in different Eastern Slovak climatic regions; Košice lies in the southern warm climatic region near the Slovak-Hungarian border, while Spišská Nová Ves is more northern in moderately warm climatic region (LAPIN et al. 2002). Phytogeographically (FUTÁK 1980), Košice is part of the Košická kotlina district (Eupannonicum, Pannonicum) and Spišská Nová Ves is in the Spišské kotliny district (Intracarpaticum, Carpaticum occidentale). Flora at these stations has been studied both recently and in the past.

Characteristics of the individual localities: the size and locality of the four railway stations are illustrated in Fig. 1 and Tab. 1; where station size is determined by the number of rails in the station (www.vlaky.net).

(1) Košice: railway station, altitude: 205 m a.s.l.; number of square and subsquare on the Slovak flora grid map (JASIČOVÁ & ZAHRADNÍKOVÁ 1976): 7293d; visited 1968–2013 (by Jehlík: 31.7.1968 and 13.7.1974; by Jehlík, Zaliberová and Májeková: 25.8.2012, 30.4.2013 and 16.7.2013).

Košice is an important rail junction with the following lines; number 180 to the North-west (Žilina – Košice); 190 to the East (Košice – Čierna nad Tisou); 160 to the West (Zvolen – Košice); 169 South to Hungary (Košice – Hidasnémeti) and 188 North to Poland (Košice – Plaveč – Muszyna). The railway station began operation in 1871 and the lines were opened between 1860 and 1890 (KUBÁČEK et al. 2013).

(2) Spišská Nová Ves: railway station, 465 m a.s.l; 7089a; visited in 1973–2013 (by Jehlík: 12.7.1973; by Jehlík, Zaliberová and Májeková: 2.5.2013, 17.7.2013).

Station lies on the Žilina – Košice 180 line and rail traffic began here in 1871. Passenger traffic on a local Spišská Nová Ves – Levoča 186 line commenced in 1892, was stopped in 2003 (KUBÁČEK et al. 2013).

(3) Plaveč station lies in North-eastern Slovakia in the moderately warm climatic region. It serves as a passenger and freight station, and connects to the 185 Poprad-Tatry and 188 Košice lines; this line previously continued into Poland (cf. MÁJEKOVÁ & LIMÁNEK 2016).

(4) Čierna nad Tisou lies in the warm climatic region of South-eastern Slovakia. It is the largest and the most important Slovak railway trans-shipment yard and also serves as a passenger railway station. The station is connected to Košice line 190 and to the Ukrainian rail network (cf. JEHLÍK et al. in press).




Fig. 1. Map of railway network in Slovakia with studied stations.

Tab. 1. Comparison of studied railway stations' size, the number of recorded species and available values (% of all taxa) in Ellenberg and Borhidi species temperature values.

Railway station	Number of rails	Number of recorded species	Ellenberg	Borhidi
Košice	119	207	148 (71%)	193 (93%)
Spišská Nová Ves	38	204	143 (70%)	191 (94%)
Plaveč	11	202	141 (70%)	189 (94%)
Čierna nad Tisou	173	582	383 (66%)	481 (83%)

Field research

Previous species lists from Košice and Spišská Nová Ves were incomplete, with only species presence/absence documented, therefore we recorded all vascular plant species identified at these railway stations; with abundance on a 5-degree scale: a – rare, b – occasional, c – scattered, d – frequent and e – abundant (cf. JEHLÍK 2013, JEHLÍK et al. in press).

Plant taxa nomenclature follows MARHOLD (1998) or is cited with the author's name, particular IUCN threat categories are according to ELIÁŠ jun. et al. (2015) and their origin and invasiveness are as in MEDVECKÁ et al. (2012). Specimens are deposited in the herbarium at the Institute of Botany, Slovak Academy of

Sciences in Bratislava (SAV) and in the herbarium of the Institute of Botany, Academy of Sciences of the Czech Republic in Průhonice (PRA).

Data analysis

Comparison of species' temperature preferences was from Košice, Spišská Nová Ves, Plaveč (MÁJEKOVÁ & LIMÁNEK 2016) and Čierna nad Tisou (JEHLÍK et al. in press) species lists. Ellenberg and Borhidi temperature indicator values were used and compared (ELLENBERG et al. 1992, BORHIDI 1995), and species' temperature values at the stations were presented as Box-Whisker plots in Statistica program.

Results and discussion

Košice

A total of 207 plant taxa were recorded at Košice railway station (Tab. 2); comprising 192 current taxa and 72 from past records; and featuring 97 native taxa and 99 alien with 10 uncertain residence status. Archaeophytes (with 63 taxa) were more abundant than neophytes (36). Plants belonged to 46 families, the most abundant were: Asteraceae (28 taxa), Poaceae (22), Brassicaceae (15), Cichoriaceae (14) and Fabaceae (10).

While *Lactuca saligna* (VU), *Chenopodium murale*, *Tribulus terrestris* (NT) and *Draba nemorosa* (LC) are recently recorded threatened species, comparison between past and recent species lists is impossible because of incomplete previous monitoring.

The discovery of *Tribulus terrestris* at Košice station documents a new distribution locality because it had only previously been recorded in dry natural and synanthropic habitats in south-western Slovakia (ZAHRADNÍKOVÁ 1982, ELIÁŠ & FERÁKOVÁ 1999). Nowadays, species is spreading northwards by railway traffic to the secondary habitats, specifically railway stations (ELIÁŠ 1977, MÁJEKOVÁ et al. 2014). Košice is beside the most eastern Čierna nad Tisou station the second species location in Eastern Slovakia (JEHLÍK et al. in press).

In addition, although *Draba nemorosa* is generally spread throughout Slovakia, our finding is the first in the Košická kotlina phytogeographical district (cf. PENIAŠTEKOVÁ & KLIMENT 2002, MÁJEKOVÁ et al. 2014).

Spišská Nová Ves

A total of 204 taxa were recorded at the Spišská Nová Ves railway station (Tab. 2); 56 from past lists and 191 recent additions. The 112 native plant taxa prevailed over 83 aliens, and 58 archaeophytes over 25 neophytes, with an additional 9 species having uncertain residence status.

Species are divided into 42 plant families, with the following most abundant: Asteraceae (24), Poaceae (24), Brassicaceae (22), Fabaceae (13) and Rosaceae (12); while threatened species comprised the recently recorded *Papaver dubium* subsp. *austromoravicum* (NT) and *Draba nemorosa* (LC).

Tab. 2. List of recorded taxa at Košice and Spišská Nová Ves railway stations.
 Historical data comes from Jehlík (1968–1974) and recent data from Jehlík, Zaliberová and Májeková (2012–2013). Annotations: x = taxa presence and a–e = taxa abundance.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Acer campestre</i>		a		
<i>Acer platanoides</i>		a		a
<i>Acetosa thrysiflora</i>	x	c		b
<i>Acinos arvensis</i>		a		
<i>Acosta rhenana</i>	x	b		
<i>Aegopodium podagraria</i>				a
<i>Aethusa cynapium</i> subsp. <i>cynapioides</i>		a		
<i>Achillea millefolium</i> agg.	x	c		c
<i>Ailanthus altissima</i>		a		
<i>Alliaria petiolata</i>				a
<i>Amaranthus albus</i>	x	a-b	x	
<i>Amaranthus powelli</i>		c		
<i>Amaranthus retroflexus</i>	x	c-d	x	c
<i>Ambrosia artemisiifolia</i>		b-c		
<i>Anchusa officinalis</i>	x		x	b
<i>Anthriscus cerefolium</i>		a		
<i>Anthriscus sylvestris</i>				b-c
<i>Apera spica-venti</i>			x	
<i>Arabidopsis thaliana</i>		b		c
<i>Arctium lappa</i>	x	b		b
<i>Arctium minus</i>	x			
<i>Arctium tomentosum</i>		a-b	x	b
<i>Arenaria serpyllifolia</i>		c		a-c
<i>Armoracia rusticana</i>			x	a
<i>Arrhenatherum elatius</i>		d		c-d
<i>Artemisia absinthium</i>	x	a-b		a
<i>Artemisia vulgaris</i>	x	c	x	c
<i>Astragalus glycyphyllos</i>				a
<i>Atriplex patula</i>		a-b		a
<i>Atriplex tatarica</i>	x	a-c		
<i>Ballota nigra</i> subsp. <i>nigra</i>		b		a-b
<i>Barbarea vulgaris</i>				a
<i>Bassia scoparia</i> subsp. <i>scoparia</i>	x	c		
<i>Berteroia incana</i>		a-b		b
<i>Betula pendula</i>		a		
<i>Bidens frondosa</i>		a-c		a
<i>Brassica napus</i> subsp. <i>napus</i>	x	b-c		a-b
<i>Bromus hordeaceus</i>		b		a-b
<i>Bromus inermis</i>	x		x	
<i>Bromus japonicus</i>		b		
<i>Bromus sterilis</i>		a		b
<i>Bromus tectorum</i>	x	a-c	x	b-c

Tab. 2. – cont.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Bunias orientalis</i>			x	a
<i>Calamagrostis epigejos</i>	x	b-c	x	c
<i>Calystegia sepium</i>				b
<i>Camelina microcarpa</i>		a		
<i>Camelina microcarpa</i> subsp. <i>sylvestris</i>			x	
<i>Capsella bursa-pastoris</i>		b-c		c
<i>Cardaminopsis arenosa</i>				a
<i>Cardaria draba</i>			x	a
<i>Carduus acanthoides</i>	x	b	x	b
<i>Carex hirta</i>	x			
<i>Carex spicata</i>			x	
<i>Cerastium glutinosum</i>				a
<i>Cerastium holosteoides</i>		b		b
<i>Cerastium semidecandrum</i>		c		c
<i>Cerasus avium</i>				a
<i>Chaerophyllum aromaticum</i>				a
<i>Chamerion angustifolium</i>			x	a
<i>Chelidonium majus</i>		a-b		a-b
<i>Chenopodium album</i>	x	b		b
<i>Chenopodium glaucum</i>	x	b		
<i>Chenopodium murale</i>		a		
<i>Chenopodium pedunculare</i> Bertol.				a
<i>Chenopodium strictum</i>		d		
<i>Chorispora tenella</i>				a
<i>Cichorium intybus</i>	x	a-c	x	c
<i>Cirsium arvense</i>	x	b-c		b
<i>Cirsium vulgare</i> subsp. <i>vulgare</i>	x			a
<i>Citrullus lanatus</i>		a		
<i>Consolida regalis</i>	x	a	x	a
<i>Convolvulus arvensis</i>	x	c-d		b-c
<i>Conyza canadensis</i>	x	b-c	x	b-c
<i>Crepis</i> sp.		b		
<i>Crepis tectorum</i>		a		a
<i>Cruciata glabra</i>				a
<i>Dactylis glomerata</i>		a		a
<i>Dalanum angustifolium</i>		a		
<i>Daucus carota</i>	x	c		a-c
<i>Descurainia sophia</i>			x	
<i>Digitaria sanguinalis</i>		c-d		c
<i>Draba nemorosa</i>		a		c
<i>Echinochloa crus-galli</i>		b		a
<i>Echium vulgare</i>	x	b-c	x	c
<i>Elytrigia repens</i>	x	c		c
<i>Epilobium collinum</i>		a		
<i>Epilobium hirsutum</i>		a		a
<i>Epilobium tetragonum</i>		a		a

Tab. 2. – cont.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Equisetum arvense</i>	x	c		b-c
<i>Eragrostis minor</i>	x	c	x	b
<i>Erigeron acris</i>		a		
<i>Erodium cicutarium</i>		a	x	
<i>Erophila verna</i>		b		b
<i>Erysimum cheiranthoides</i>				a
<i>Falcaria vulgaris</i>	x			
<i>Fallopia convolvulus</i>		c		c
<i>Festuca pratensis</i>				a
<i>Festuca rubra</i>		b		b
<i>Fragaria vesca</i>		a		a
<i>Galinsoga parviflora</i>	x	a-c	x	a
<i>Galinsoga urticifolia</i>		a-b		
<i>Galium album</i>		b-c		b-c
<i>Galium aparine</i>		b		
<i>Galium odoratum</i>				a
<i>Galium spurium</i>		a		
<i>Galium verum</i>	x	b		a
<i>Geranium pratense</i>	x		x	b
<i>Geranium purpureum</i> Vill.		a		
<i>Geranium pusillum</i>		a-b		
<i>Geranium robertianum</i>		b		b
<i>Geranium sibiricum</i>		a		a-c
<i>Glechoma hederacea</i>				b
<i>Helianthus annuus</i> var. <i>macrocarpus</i> (DC.) Cockerell	x			a
<i>Heracleum sphondylium</i>				b
<i>Hieracium sabaudum</i>		a		
<i>Hordeum murinum</i>		a		a
<i>Humulus lupulus</i>		a		
<i>Hypericum perforatum</i>		b-c		a-b
<i>Impatiens parviflora</i>		b		c
<i>Iva xanthiifolia</i>	x	a		
<i>Juglans regia</i>		a		
<i>Lactuca saligna</i>		b		
<i>Lactuca serriola</i>	x	b-c	x	b-c
<i>Lamium album</i>		a		b-c
<i>Lamium amplexicaule</i>		b		a
<i>Lamium purpureum</i>		a		
<i>Lappula squarrosa</i>				
<i>Lapsana communis</i>				a
<i>Lathyrus pratensis</i>				a
<i>Lathyrus tuberosus</i>	x	a	x	a
<i>Leontodon autumnalis</i>		a-b		a
<i>Lepidium campestre</i>	x		x	
<i>Lepidium densiflorum</i> (incl. <i>L. neglectum</i>)	x	b	x	a

Tab. 2. – cont.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Lepidium ruderale</i>	x	b	x	b
<i>Libanotis pyrenaica</i>		a	x	c
<i>Linaria vulgaris</i>	x	a-c		c
<i>Lithospermum arvense</i>		b		a
<i>Lolium perenne</i>	x	a-c		c
<i>Lotus corniculatus</i>	x	b-c		
<i>Lycopersicon esculentum</i>		a		
<i>Malus domestica</i>	x	b		a
<i>Malva neglecta</i>				a
<i>Matricaria discoidea</i>		a	x	b
<i>Medicago falcata</i>			x	a
<i>Medicago lupulina</i>		c		c
<i>Medicago sativa</i>	x	a-c		
<i>Medicago × varia</i>		a-b		b
<i>Melilotus albus</i>	x	a-b	x	a
<i>Melilotus officinalis</i>	x	b-c	x	b
<i>Microrrhinum minus</i>	x	a-c		a
<i>Myosotis arvensis</i>				b
<i>Myosotis sparsiflora</i>				a
<i>Myosotis stricta</i>		c		b
<i>Negundo aceroides</i>		b-c		
<i>Oenothera biennis</i>	x	a-b		a
<i>Oenothera rubricaulis</i>		a		
<i>Oenothera villosa</i>	x	b	x	a
<i>Onopordum acanthium</i>	x		x	a
<i>Origanum vulgare</i> subsp. <i>vulgare</i>				a
<i>Panicum capillare</i> subsp. <i>barbipulvinatum</i> (Nash) Tzvel.		c		
<i>Papaver dubium</i> subsp. <i>austromoravicum</i>				a
<i>Papaver rhoeas</i>		b		a
<i>Parthenocissus inserta</i> (A. Kern.) Fritsch		a		a
<i>Pastinaca sativa</i> subsp. <i>sativa</i>	x	b-d	x	a-b
<i>Persica vulgaris</i>	x			
<i>Persicaria lapathifolia</i>			x	
<i>Persicaria maculosa</i>	x	a-b		b
<i>Picris hieracioides</i>	x	b-c	x	b-c
<i>Pilosella officinarum</i>		a-b		a
<i>Pimpinella saxifraga</i>		b		a
<i>Pinus sylvestris</i>				a
<i>Plantago lanceolata</i>		c		c
<i>Plantago major</i>	x	b-c		b
<i>Plantago media</i>				a
<i>Poa annua</i>		a		b
<i>Poa bulbosa</i> subsp. <i>vivipara</i>				a
<i>Poa compressa</i>		a-b	x	a-b
<i>Poa palustris</i>				a
<i>Poa pratensis</i>		d		c

Tab. 2. – cont.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Polygonum aviculare</i> agg.	x	d		c
<i>Populus alba</i>		a		
<i>Portulaca oleracea</i>		b-c		b
<i>Potentilla anserina</i>				a
<i>Potentilla argentea</i>		b-c	x	b-c
<i>Potentilla intermedia</i>				b-c
<i>Potentilla recta</i>			x	a
<i>Potentilla reptans</i>		c		b
<i>Potentilla supina</i>			x	a
<i>Prenanthes purpurea</i>				a
<i>Puccinellia distans</i>	x	a-b	x	a
<i>Ranunculus repens</i>				a
<i>Reseda lutea</i>	x	a-b	x	a
<i>Rorippa sylvestris</i>		a		
<i>Rosa canina</i>		a-b		a-b
<i>Rubus caesius</i>		b-c		b-c
<i>Rubus idaeus</i>		a		a
<i>Rumex confertus</i>	x			
<i>Rumex crispus</i>	x			a
<i>Rumex obtusifolius</i>		b		
<i>Rumex patientia</i>		a		
<i>Salix caprea</i>		b		a-b
<i>Salix cinerea</i>		a		a
<i>Salix silesiaca</i>				a
<i>Salvia verticillata</i>		a	x	a
<i>Sambucus nigra</i>	x	a-c		a-c
<i>Saponaria officinalis</i>	x	a-c	x	a-b
<i>Scrophularia nodosa</i>		a		
<i>Securigera varia</i>		a	x	b
<i>Sedum acre</i>		b		b
<i>Senecio ovatus</i>				a
<i>Senecio vernalis</i>		b		b
<i>Senecio viscosus</i>	x	a-b		a
<i>Senecio vulgaris</i>		b		a-b
<i>Setaria verticillata</i>	x	a		
<i>Setaria viridis</i> (L.) P. B. subsp. <i>pycnocoma</i> (Steud.) Tzvelev				a
<i>Setaria viridis</i> subsp. <i>viridis</i>		c-d		b
<i>Silene latifolia</i> subsp. <i>alba</i>		b-c		b-c
<i>Silene noctiflora</i>		a		
<i>Silene vulgaris</i>		a		a
<i>Sisymbrium altissimum</i>	x		x	
<i>Sisymbrium loeselii</i>		a	x	
<i>Sisymbrium officinale</i>			x	
<i>Solanum nigrum</i>		b		
<i>Solanum nigrum</i> subsp. <i>schultesii</i>		a		

Tab. 2. – cont.

Taxon	Košice		Spišská Nová Ves	
	historic	recent	historic	recent
<i>Solanum tuberosum</i>			x	
<i>Solidago canadensis</i>		b-c		b
<i>Solidago gigantea</i>				a
<i>Sonchus arvensis</i>	x			
<i>Sonchus asper</i>		a		
<i>Sonchus oleraceus</i>		b-c		c
<i>Stachys annua</i>		b		
<i>Stellaria graminea</i>		a		
<i>Stellaria media</i>		c		b
<i>Stenactis annua</i>		a-c		b
<i>Swida sanguinea</i>		a-b		a
<i>Symphytum officinale</i>				a
<i>Tanacetum vulgare</i>	x	a-c	x	a-b
<i>Taraxacum sect. Ruderalia</i>		b-d		b-d
<i>Thlaspi arvense</i>		b		
<i>Thlaspi perfoliatum</i>		a		
<i>Tilia cordata</i>				a
<i>Tilia platyphyllos</i>		a		a
<i>Tithymalus cyparissias</i>		b		a
<i>Tithymalus helioscopia</i>				b
<i>Torilis japonica</i>		b		
<i>Tragopogon dubius</i>		c		
<i>Tragopogon orientalis</i>		c		
<i>Tribulus terrestris</i>		a-b		
<i>Trifolium pratense</i>		c		
<i>Trifolium repens</i>		b-d		b-c
<i>Tripleurospermum perforatum</i>	x	b-c	x	b-c
<i>Tussilago farfara</i>	x	a		a
<i>Ulmus glabra</i>				a
<i>Urtica dioica</i>		b-c		b
<i>Valerianella locusta</i>		b		a
<i>Verbascum blattaria</i>		a		
<i>Verbascum lychnitis</i>			x	a-b
<i>Veronica arvensis</i>		b		b
<i>Veronica hederifolia</i>		b		a
<i>Veronica persica</i>				a
<i>Veronica polita</i>				b
<i>Vicia cracca</i>				b
<i>Vicia villosa</i>				a
<i>Viola arvensis</i>		a-b		b-c
<i>Viola cf. odorata</i>		b		
<i>Viola odorata</i>				b
<i>Viola tricolor</i>		a		a
<i>Vitis vinifera</i>		a		
<i>Xanthoxalis stricta</i>		a		

Comparison of the species structure

Tab. 1 highlights that although Košice's 119 tracks makes it a much larger railway complex than Spišská Nová Ves with its 38 tracks, the number of recorded species is quite similar at 207 compared to 204; and 192 to 191 in recent listing. Specifically, there is uniformity of occurrence of the following large group of thermophilous species at these stations: *Amaranthus albus*, *A. retroflexus*, *Anchusa officinalis*, *Artemisia absinthium*, *Bromus tectorum*, *Consolida regalis*, *Conyza canadensis*, *Digitaria sanguinalis*, *Echium vulgare*, *Eragrostis minor*, *Onopordum acanthium*, *Portulaca oleracea*, *Setaria viridis*, *Sisymbrium altissimum* and *S. loeselii*. However, differences in populations were also noted; where the following thermophilous species were recorded only at Košice: *Ambrosia artemisiifolia*, *Amaranthus powellii*, *Atriplex tatarica*, *Geranium purpureum*, *Iva xanthiifolia*, *Lactuca saligna*, *Lappula squarrosa*, *Rumex patientia*, *Setaria verticillata* and *Tribulus terrestris*.

The most novel finding was *Chorispora tenella*'s presence at Spišská Nová Ves station. This is not only a new locality for this species but also its northernmost sighting in Slovakia; as it was previously recorded mainly at a few South-western and eastern Slovak locations (MICHALKOVÁ 2002). Its presence at Spišská Nová Ves can be explained as introduction by railway traffic from southern warmer areas. The species was recorded, for example, in Dobrá and Čierna nad Tisou South-eastern Slovak railyards (JEHLÍK et al. in press).

Archaeophytes prevailed over neophytes at both Košice and Spišská Nová Ves railway stations; supporting CHYTRÝ et al.'s (2008) finding that archaeophytes have higher affinity for open vegetation at dry or mesic sites and neophytes prefer wet habitats and woodlands. Representation of dominant plant families at these two sites is also similar to that at south-eastern stations (JEHLÍK et al. in press).

We employed the different Ellenberg and Borhidi indicator scales to evaluate species temperature preferences, and we also included the further two Čierna nad Tisou and Plaveč railway stations to supplement Košice and Spišská Nová Ves results. Many more species were recorded at the southern-most Čierna nad Tisou station because this is the largest rail area with the greatest railway transhipment yard in Slovakia, and therefore it has the highest diaspore introduction rates (Tab.1); moreover this area was researching for the longest time. We found the same median value (6) on the thermophilous scale at all four stations, and Ellenberg and Borhidi indicator values returned similar results (Fig. 2). While Košice and Spišská Nová Ves have the same value range and Plaveč is very similar with only small deviation in the lower values, the Čierna nad Tisou value range is higher.

While our research shows similar thermophilous species representation at the four studied railway stations and climate has no distinguishing role, the heated rail and gravel substrate in these railway vicinities provide a fruitful location for thermophilous species which do not thrive so well in less suitable temperatures at native biotopes in colder areas.




Fig. 2. Box and Whiskers plots of species temperature preferences by a) Ellenberg, b) Borhidi scales at studied railway stations (1 – Košice, 2 – Spišská Nová Ves, 3 – Plaveč, 4 – Čierna nad Tisou).

Use of environmental indicator values

Our research revealed differences in practical use of the indicator scales. The Ellenberg scale was proposed for Central Europe and most often used in analyses. However, this scale determined more species with missing values in our dataset (Tab. 1) than the Borhidi indicator values proposed for Hungary and whose proximity to our study area proved more reliable for our data. Since this differentiation is supported by KOCHJAROVÁ et al.'s (2015) results, it is advisable to test these scales before analysis.

Acknowledgments

We are grateful to Slovak Republic Railways for permission for this study and we thank Iveta Škodová for reading and manuscript comments and Raymond Marshall for English revision.

References

- BORHIDI A. (1995): Social behaviour types, the naturalness and relative ecological indicator values of the higher plants in the Hungarian Flora. – Act. Bot. Hung. 39: 97-181.
- BRANDES D. (1983): Flora und Vegetation der Bahnhöfe Mitteleuropas. – Phytocoenologia 11: 31-115.
- BRANDES D. (2008): Bibliographie zur Eisenbahnvegetation, www.digibib.tu-bs.de/?docid=00021885 (accessed 20-July-2016)
- CHYTRY M., MASKELL L. C., PINO J., PYŠEK P., VILÀ M., FONT X. & SMART S. M. (2008): Habitat invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental and oceanic regions of Europe. – J. Appl. Ecol. 45: 448-458.
- ELIÁŠ P. (1977): Poznámky k lokalitám *Tribulus terrestris* L. subsp. *orientalis* (Kern.) Dost. na železničných stanovištiach Slovenska. – Zprávy Českoslov. Bot. Společn. 12: 127-129.

- ELIÁŠ P. (1979a): Zriedkavejšie rastliny železničných komunikácií na západnom Slovensku I. – Biológia 34: 67-70.
- ELIÁŠ P. (1979b): *Linario-Brometum tectorum* na železničnej stanici Cífer (západné Slovensko). – Biológia 34: 329-333.
- ELIÁŠ P. (1979c): The association *Conzyo-Cynodontetum dactyloni* in western Slovakia, Czechoslovakia. – Preslia 51: 349-362.
- ELIÁŠ P. (1981): Zriedkavejšie rastliny železničných komunikácií na západnom Slovensku II. – Biológia 36: 73-77.
- ELIÁŠ P. & FERÁKOVÁ V. (1999): *Tribulus terrestris* L. – In: Čeřovský J., Feráková V., Holub J., Maglocký Š. & Procházka F.: Červená kniha ohrozených a vzácných druhov rastlín a živočíchov SR a ČR, Vol. 5. Vyššie rastliny, p. 382. – Príroda, Bratislava.
- ELIÁŠ P. jun. (2011): *Geranium purpureum* Vill. – new alien species to the Slovak flora. – Thaiszia 21: 21-28.
- ELIÁŠ P. jun., DÍTĚ D., ELIAŠOVÁ M. & ĎURIŠOVÁ L. (2013): Distribution and origin of *Aegilops* species in Slovakia. – Thaiszia 23: 117-129.
- ELIÁŠ P. jun., DÍTĚ D., KLIMENT J., HRIVNÁK R. & FERÁKOVÁ V. (2015): Red list of ferns and flowering plants of Slovakia, 5th edition (October 2014). – Biológia 70: 218-228.
- ELLENBERG H., WEBER H. E., DÜLL R., WIRTH V., WERNER W. & PAULISSEN D. (1992): Zeiger-werte von Pflanzen in Mitteleuropa. 2nd ed. – Scr. Geobot. 18: 1-258.
- FRANTOVA J. (1947): Plevelová, ruderálna a adventívna flóra okolia Trnavy. – Prírod. Sborn. 2: 153-248.
- FUTÁK J. (1980): Fytogeografické členenie. – In: MAZÚR E. (ed.): Atlas Slovenskej socialistickej republiky, p. 88. – Slovenská akadémia vied, Slovenský úrad geodézie a kartografie, Bratislava.
- GALERA H., SUDNIK-WÓJCIKOWSKA B., WIERZBICKA M. & WIŁKOMIRSKI B. (2012): Directions of changes in the flora structure in the abandoned railway areas. – Ecol. Quest. 16: 29-39.
- GALERA H., SUDNIK-WÓJCIKOWSKA B., WIERZBICKA M., JARZYNA I. & WIŁKOMIRSKI B. (2014): Structure of the flora of railway areas under various kinds of anthropopression. – Polish Bot. J. 59: 1-10.
- HOHLA M., KLEESADL G. & MELZER H. (2000): Neues zur Flora des oberösterreichischen Bahnanlagen – mit Einbeziehung einiger grenznaher Bahnhöfe Bayerns. – Beitr. Naturk. Oberösterreichs 9: 191-250.
- HOHLA M., KLEESADL G. & MELZER H. (2002): Neues zur Flora der oberösterreichischen Bahnanlagen – mit Einbeziehung einiger Bahnhöfe Bayerns – Fortsetzung. – Beitr. Naturk. Oberösterreichs 11: 507-578.
- HOHLA M., KLEESADL G. & MELZER H. (2005): Neues zur Flora der oberösterreichischen Bahnanlagen. – Beitr. Naturk. Oberösterreichs 14: 147-199.
- HOSKOVEC L. (2009): Květena pražských nádraží: Nejčastejší druhy rostlin železničných nádraží Prahy, www.botany.cz/cs/soupis-rostlin-prazskych-nadrazi (accessed 20-July-2016)
- HUBER W. (1992): Zur Ausbreitung von Blütenpflanzenarten an Sekundärstandorten der Nordschweiz. – Bot. Helv. 102: 93-108.
- JASIČOVÁ M. & ZAHRADNÍKOVÁ K. (1976): Organizácia a metodika mapovania rozšírenia rastlinných druhov v západnej treťine Slovenska. – Biológia 31: 74-80.
- JEHLÍK V. (1986): The vegetation of railways in Northern Bohemia (eastern part). – Vegetace ČSSR, Ser. A, 14: 1-368.
- JEHLÍK V. (1995): Současný výskyt cizích expanzívnych druhov rostlin v železničných uzlech České republiky. – Ochr. Rostl. 31: 149-160.
- JEHLÍK V. (ed.) (1998): Cizí expanzívní plevely České republiky a Slovenské republiky. – Academia, Praha, 508 pp.

- JEHLÍK V. (2013): Die Vegetation und Flora der Flusshäfen Mitteleuropas. – Academia, Praha, 546 pp.
- JEHLÍK V. & DOSTÁLEK J. (1989): *Convolvulo arvensis-Botriochloaetum ischaemi* – eine neue Apophytengesellschaft auf Eisenbahnhöfen in der Südslowakei. – Preslia 61: 43-50.
- JEHLÍK J. & DOSTÁLEK J. (2008): Influence of railway transport in the South-East of Slovakia on formation of adventive flora in Central Europe. – Biodiv. Res. Conserv. 11-12: 27-32.
- JEHLÍK V., MÁJEKOVÁ J. & ZALIBEROVÁ M. (2013): New discovered adventive plants from eastern Slovakia. – Thaiszia 23: 61-66.
- JEHLÍK V., ZALIBEROVÁ M. & MÁJEKOVÁ J. (in press): Influence of the Eastern migration route on Slovak flora – comparison after 40 years. – Tuexenia.
- KOCHJAROVÁ J., ŠKODOVÁ I. & BLANÁR D. (2015): Grasslands in the border area of Carpathian and Pannonian regions: an example from Muránska planina Mts (Central Slovakia). – Tuexenia 35: 195-220.
- KUBÁČEK J. et al. (2013): Dejiny železníc na území Slovenska. – Železnice Slovenskej republiky, Bratislava, 280 pp.
- LAPIN M., FAŠKO P., MELO M., ŠŤASTNÝ P. & TOMLAIN J. (2002): Klimatické oblasti. – In: MIKLÓS L. (ed.): Atlas krajiny Slovenskej republiky, p. 95. – Ministerstvo životného prostredia, Slovenská agentúra životného prostredia, Bratislava, Banská Bystrica.
- MÁJEKOVÁ J., ZALIBEROVÁ M. & JEHLÍK V. (2013): Extinct species *Ceratocephala testiculata* (Crantz) Besser rediscovered in Slovakia after 44 years. – Thaiszia 23: 141-145.
- MÁJEKOVÁ J., LETZ D. R., SLEZÁK M., ZALIBEROVÁ M. & HRIVNÁK R. (2014): Rare and threatened vascular plants of the railways in Slovakia. – Biodiv. Res. Conserv. 35: 75-85.
- MÁJEKOVÁ J. & LIMÁNEK J. (2016): Diverzita flóry železničných staníc na trati Poprad – Plaveč (východné Slovensko). – Bull. Slov. Bot. Spoločn., 38/1: 37-45.
- MARHOLD K. (ed.) (1998): Papraďorasty a semenné rastliny. – In: MARHOLD K. & HINDÁK F. (eds.): Zoznam nižších a vyšších rastlín Slovenska, p. 333-687. – Veda, Bratislava.
- MEDVECKÁ J., KLIMENT J., MÁJEKOVÁ J., HALADA L., ZALIBEROVÁ M., GOJDÍCOVÁ E., FERÁKOVÁ V. & JAROLÍMEK I. (2012): Inventory of the alien flora of Slovakia. – Preslia 84: 257-309.
- MICHALKOVÁ E. (2002): *Chorispora* R. Br. ex DC. – In: GOLIAŠOVÁ K. & ŠÍPOŠOVÁ H. (eds.): Flóra Slovenska V/4, p. 260-264. – Veda, Bratislava.
- NOWINSKA R. & CZARNA A. (2008): Impact of Railway Facility Operation on Floral Growth in Powodowo, the Region of Wielkopolska. – Polish J. of Environ. Stud. 17: 613-622.
- PENIAŠTEKOVÁ M. & KLIMENT J. (2002): *Draba* L. – In: Goliašová K. & Šípošová H. (eds.): Flóra Slovenska V/4, p. 500-540. – Veda, Bratislava.
- PISKORZ R. & CZARNA A. (2006): Vascular plants on active and closed railway stations in Wolsztyn and its surroundings. – Roczniki Akademii Rolniczej w Poznaniu, Botanica Ser. Steciana, 10: 137-156.
- RABITSCH W. & ESSL F. (2006): Biological invasions in Austria: patterns and case studies. – Biol. Invas. 8: 295-308.
- RŮŽIČKA V. & KOBLÍŽEK J. (2009): Kakost nachový (*Geranium purpureum*), nový druh pro květenu České republiky. – Zprávy Čes. Bot. Společ. 44: 23-27.
- SCHIDLAY E. (1944): Šírenie sa rastlín *Erucastrum nasturtiifolium* a *E. gallicum* na Slovensku. (Príspevok k flóre Slovenských železníc). – Techn. Obzor Slov., Prírod. Príl.: 49-55.
- SHEVERA M. V., MÁJEKOVÁ J., ZALIBEROVÁ M., PROTOPOPOVA V. V. & ANDRIK E. J. (2015): *Geranium purpureum* (Geraniaceae), a new alien species of the flora of Ukrainian plain area. – Ukr. Bot. J. 72: 334-339.

- STAŘECKÁ C. (2010): Diverzita flóry železničních tratí ve středních Čechách. – Studia Oecologica 4: 70-76.
- VAN KLEUNEN M. et al. (2015): Global exchange and accumulation of non-native plants. – Nature 525: 100-103.
- WIĘKOMIRSKI B., GALERA H., SUDNIK-WÓJCIKOWSKA B., STASZEWSKI T. & MALAWSKA M. (2012): Railway Tracks – Habitat Conditions, Contamination, Floristic Settlement – A Review. – Environment and Natural Resources Research 2(1): 86-95.
- ZALIBEROVÁ M. & MAJEKOVÁ J. (2014): Poznámky k prvému nálezu *Geranium purpureum* Vill. na Slovensku a rozšírenie druhu na železničných staniciach Záhorie (západné Slovensko). – Bull. Slov. Bot. Spoločn. 36(2): 221-230.
- ZAHRADNÍKOVÁ K. (1982): *Zygophyllaceae*. – In: Futák J. & Bertová L. (eds.): Flóra Slovenska III, p. 535-537. – Veda, Bratislava.

Received: October 24th 2016
Revised: December 02nd 2016
Accepted: December 02nd 2016