



P. J. ŠAFÁRIK UNIVERSITY
FACULTY OF SCIENCE
INSTITUTE OF MATHEMATICS
Jesenná 5, 041 54 Košice, Slovakia



S. Cichacz and M. Horňák

**Decomposition of bipartite graphs
into closed trails**

IM Preprint, series A, No. 2/2007
February 2007

Decomposition of bipartite graphs into closed trails*

Sylvia Cichacz

AGH University of Science and Technology

Al. Mickiewicza 30, 30-059 Kraków, Poland, cichacz@agh.edu.pl

Mirko Hornák

P.J. Šafárik University

Jesenná 5, 040 01 Košice, Slovakia, mirko.hornak@upjs.sk

Abstract. Let $\text{Lct}(G)$ denote the set of all lengths of closed trails that exist in an even graph G . A sequence (t_1, \dots, t_p) of terms of $\text{Lct}(G)$ adding up to $|E(G)|$ is G -realizable provided there is a sequence (T_1, \dots, T_p) of pairwise edge-disjoint closed trails in G such that T_i is of length t_i for $i = 1, \dots, p$. The graph G is arbitrarily decomposable into closed trails if all possible sequences are G -realizable. In the paper it is proved that if $a \geq 1$ is an odd integer and $M_{a,a}$ is a perfect matching in $K_{a,a}$, then the graph $K_{a,a} - M_{a,a}$ is arbitrarily decomposable into closed trails.

Keywords: even graph, closed trail, graph arbitrarily decomposable into closed trails, bipartite graph

MSC 2000: 05C70

All graphs we are dealing with in this paper are simple, finite and nonoriented. We use the standard terminology and notation of graph theory.

For $p, q \in \mathbb{Z}$ let $[p, q]$ denote the *integer interval* bounded by p and q , i.e. $[p, q] := \{z \in \mathbb{Z} : p \leq z \leq q\}$; similarly, let $[p, \infty) := \{z \in \mathbb{Z} : p \leq z\}$. The *concatenation* of finite sequences $A = (a_1, \dots, a_m)$ and $B = (b_1, \dots, b_n)$ is the sequence $AB := (a_1, \dots, a_m, b_1, \dots, b_n)$. The concatenation is an associative operation on finite sequences; we use this fact in the notation $\prod_{i=1}^k A_i$ representing the concatenation of finite sequences A_i , $i \in [1, k]$, in the order given by the sequence (A_1, \dots, A_k) . As usual, A^k denotes $\prod_{i=1}^k A_i$ with $A_i = A$ for any $i \in [1, k]$, and A^0 is the empty sequence $()$. A finite sequence $A = (a_1, \dots, a_m)$ is

*The work of the first author was supported by Visegrad Scholarship nr S-016-2006. The work of the second author was supported by Science and Technology Assistance Agency under the contract No. APVT-20-004104 and by the Slovak grant VEGA 1/3004/06.

changeable to a finite sequence $A = (a'_1, \dots, a'_m)$ of the same length (in symbols $A \sim A'$) if there is a bijection $\pi \subseteq [1, m] \times [1, m]$ such that $a'_i = a_{\pi(i)}$ for any $i \in [1, m]$. If $I \subseteq [1, m]$, we denote by $A\langle I \rangle$ the subsequence of A formed by all a_i 's with $i \in I$ (ordered in compliance with the natural ordering of I).

A *closed trail* of length $n \in [3, \infty)$ (an *n-trail* for short) is a sequence $\prod_{i=1}^{n+1} (x_i)$ of vertices of G such that $x_1 = x_{n+1}$ and if $i, j \in [1, n]$, $i \neq j$, then $\{x_i, x_{i+1}\} \in E(G)$ and $\{x_i, x_{i+1}\} \neq \{x_j, x_{j+1}\}$. A graph G is Eulerian if it has a closed trail of length $|E(G)|$. It is well known that a graph of order at least three is Eulerian if and only if it is connected and *even* (all its vertices are of even degrees). Thus, we may identify the notions of a closed trail in a graph G and a nontrivial connected even subgraph of G . Let $\text{Lct}(G)$ be the set of all lengths of closed trails existing in G and let $\text{Sct}(G)$ be the set of all finite sequences consisting of terms of $\text{Lct}(G)$ that add up to $|E(G)|$. Deleting a closed trail from an even graph G yields an even subgraph of G . Continuing this process until all edges of G are exhausted leads to a sequence $\tilde{T} := (\tilde{T}_1, \dots, \tilde{T}_p)$ of pairwise edge-disjoint closed trails in G such that, for any $i \in [1, p]$, $\tilde{t}_i := |E(\tilde{T}_i)| \in \text{Lct}(G)$, and $\tilde{\tau} := (\tilde{t}_1, \dots, \tilde{t}_p) \in \text{Sct}(G)$; the sequence $\tilde{\tau}$ is said to be *G-realizable* and the sequence \tilde{T} is a *G-realisation* of the sequence $\tilde{\tau}$. An even graph G is *arbitrarily decomposable into closed trails* (ADCT) provided all sequences of $\text{Sct}(G)$ are *G-realizable*.

There are some classes of even graphs that are known to be ADCT. Among these are complete graphs K_n for n odd, the graphs $K_n - M_n$, where M_n is a perfect matching in K_n , for n even (Balister [1]) and complete bipartite graphs $K_{a,b}$ for a, b even (Hornák and Woźniak [8]). An even graph that is large and dense enough is necessarily ADCT. Namely, according to Balister [2], there are positive constants n and ε such that an even graph G is ADCT whenever $|V(G)| \geq n$ and $\delta(G) \geq (1 - \varepsilon)|V(G)|$. Hornák and Kocková [7] proved that if an even complete tripartite graph $K_{p,q,r}$ with $p \leq q \leq r$ is ADCT, then either $(p, q, r) \in \{(1, 1, 3), (1, 1, 5)\}$ or $p = q = r$; moreover, the graphs $K_{1,1,3}$, $K_{1,1,5}$ and $K_{p,p,p}$ with $p = 5 \cdot 2^l$, $l \in [0, \infty)$, are ADCT. The notion of an ADCT graph can be generalized in a natural way to oriented graphs (see Balister [3] and Cichacz [5]) and to pseudographs (Cichacz et al. [6]).

It may happen that an even graph is not ADCT though all its connected components are. For example, both C_8 (an 8-vertex cycle) and $K_{2,4}$ are ADCT, but $C_8 \cup K_{2,4}$ is not since the sequence $(4)^4 \in \text{Sct}(C_8 \cup K_{2,4})$ is not $(C_8 \cup K_{2,4})$ -realizable. On the other hand, if the graphs G^1, G^2 are ADCT and $E(G^1) \cap E(G^2) = \emptyset$, but $V(G^1) \cap V(G^2) \neq \emptyset$, when trying to prove that a sequence $\tau \in \text{Sct}(G^1 \cup G^2)$ is $(G^1 \cup G^2)$ -realizable, we have at our disposal not only closed trails of G^1 and G^2 , but also closed trails $T^1 \cup T^2$, where T^i is a closed trail of G^i , $i = 1, 2$, and $V(T^1) \cap V(T^2) \neq \emptyset$. Therefore, a potential general strategy for proving that a graph G is ADCT can be described as follows: Write G as an edge-disjoint, but not vertex-disjoint, union of ADCT graphs G^1 and G^2 , and require from G^i -realizations, $i = 1, 2$, to have an additional property that some of their chosen trails contain common vertices of $V(G^1) \cap V(G^2)$.

Clearly, when analyzing whether a nontrivial connected even graph G is ADCT, it is sufficient to show that any sequence $(t_1, \dots, t_p) \in \text{Sct}(G)$ of length $p \geq 2$ is G -realizable; indeed, the graph G is Eulerian, and so the unique sequence $(|E(G)|)$ of length 1 in $\text{Sct}(G)$ is trivially G -realizable. We have also the following evident statement:

Lemma 1 *If G is an even graph, $\tau_1, \tau_2 \in \text{Sct}(G)$ and $\tau_1 \sim \tau_2$, then the sequence τ_1 is G -realizable if and only if τ_2 is.* ■

Pick disjoint sets $X^j = \{x_i^j : i \in [1, \infty)\}$, $j = 1, 2$, and let $X_{p,q}^j := \{x_i^j : i \in [p, q]\}$ for $p, q \in [1, \infty)$. In this paper the complete bipartite graph $K_{a,b}$ will have the bipartition $\{X_{1,a}^1, X_{1,b}^2\}$ and $M_{a,a}$ will be the perfect matching in $K_{a,a}$ consisting of $\{x_i^1, x_i^2\}$ for $i \in [1, a]$. If a is odd, then $K'_{a,a} := K_{a,a} - M_{a,a}$ is an even graph. The main aim of our paper is to show that the graph $K'_{a,a}$ is ADCT for any odd $a \in [1, \infty)$. We proceed by induction on a and we use the above general strategy. For odd $a \geq 7$ consider the even subgraph $F_a \cong K'_{a-4, a-4}$ of $K'_{a,a}$ induced on the set $X_{5,a}^1 \cup X_{5,a}^2$. The even graph $H_a := K'_{a,a} - F_a$ is an edge-disjoint union of the even graph $K'_{5,5}$ and two even subgraphs $G_a^1 \cong G_a^2 \cong K_{4, a-5}$ of $K'_{a,a}$ where G_a^i is induced on the set $X_{1,4}^i \cup X_{6,a}^{3-i}$, $i = 1, 2$. Thus putting $G_a := K'_{5,5} \cup G_a^1$ we obtain $H_a = G_a \cup G_a^2$. We shall show subsequently that the graphs $K'_{5,5}$ and G_a, H_a are ADCT; furthermore, G_a -realizations and H_a -realizations can be chosen to have appropriate additional properties. Note that all mentioned graphs are bipartite. The following assertion shows the maximal extent of the set $\text{Lct}(G)$ for an even bipartite graph G .

Proposition 2 *If G is an even bipartite graph, then $\text{Lct}(G) \subseteq \{2k : k \in [2, |E(G)|/2 - 2]\} \cup \{|E(G)|\}$.*

Proof. All subgraphs of G are bipartite, hence all closed trails in G (as edge-disjoint unions of cycles) are of even lengths. A subgraph T of G with $|E(T)| = |E(G)| - 2$ is not even (and therefore not a closed trail) for $G - T$ has at least two vertices of degree one. ■

When proving that an even bipartite graph G is ADCT we do not exhibit the structure of $\text{Lct}(G)$ explicitly, but we show implicitly that $\text{Lct}(G)$ is of maximal extent by finding all G -realizations that are theoretically possible from the point of view of Proposition 2.

Recall again the result on complete bipartite graphs:

Theorem 3 *If a, b are even integers with $2 \leq a \leq b$, then the graph $K_{a,b}$ is ADCT.*

We know due to Chou et al. [4] that sequences of $\text{Sct}(K_{a,b})$ with small terms have $K_{a,b}$ -realizations consisting of cycles:

Theorem 4 *If a, b are even integers with $a \geq 4$, $b \geq 6$ and $\tau = (t_1, \dots, t_p) \in \text{Sct}(K_{a,b})$ with $t_i \in \{4, 6, 8\}$ for any $i \in [1, p]$, then there is a $K_{a,b}$ -realization (T_1, \dots, T_p) of the sequence τ such that T_i is a cycle for any $i \in [1, p]$. ■*

Clearly, when analyzing whether a connected even graph G is ADCT, it is sufficient to show that any sequence $(t_1, \dots, t_p) \in \text{Sct}(G)$ of length $p \geq 2$ is G -realizable; indeed, the graph G is Eulerian, and so the unique sequence $(|E(G)|)$ of length 1 in $\text{Sct}(G)$ is trivially G -realizable.

We start our analysis by dealing with $a \leq 5$.

Proposition 5 *The graph $K'_{a,a}$ with $a \in \{1, 3, 5\}$ is ADCT.*

Proof. We have $K'_{1,1} \cong 2K_1$, and so for $a = 1$ the result follows from $\text{Sct}(K'_{1,1}) = \text{Lct}(K'_{1,1}) = \emptyset$.

Since $K'_{3,3} \cong C_6$, the unique sequence $(6) \in \text{Sct}(K'_{3,3})$ is trivially $K'_{3,3}$ -realizable.




Figure 1: $K'_{5,5}$ -realizations of three sequences

The sequences $(4)^5$, $(4)^2(6)^2$ and $(6)^2(8)$ are $K'_{5,5}$ -realizable, see Figure 1. Observe that any two 4-trails of the left $K'_{5,5}$ -realization have a common vertex, hence every sequence in $\text{Sct}(K'_{5,5})$, whose all terms are divisible by 4, is $K'_{5,5}$ -realizable. Moreover, in the middle $K'_{5,5}$ -realization any 4-trail has a common vertex with any 6-trail. Therefore, the remaining sequences $(4, 6, 10)$, $(6, 14)$, $(10)^2 \in \text{Sct}(K'_{5,5})$ are $K'_{5,5}$ -realizable, too. ■

We shall need also the following three simple statements:

Proposition 6 *If G is a complete bipartite graph with bipartition $\{X, Y\}$ and $\pi \subseteq X \times X$, $\rho \subseteq Y \times Y$ are bijections, then the mapping $\alpha \subseteq V(G) \times V(G)$ with $\alpha|_X = \pi$ and $\alpha|_Y = \rho$ is an automorphism of G . ■*

Proposition 7 *If $a \in [1, \infty)$ and $\pi \subseteq [1, a] \times [1, a]$ is a bijection, then the mappings $\bar{\pi}, \tilde{\pi} \subseteq V(K'_{a,a}) \times V(K'_{a,a})$ determined by $\bar{\pi}(x_i^j) = x_{\pi(i)}^j$ and $\tilde{\pi}(x_i^j) = x_{\pi(i)}^{3-j}$ for any $i \in [1, a]$ and $j \in [1, 2]$, are automorphisms of $K'_{a,a}$. ■*

Lemma 8 *If T_1, T_2 are edge-disjoint closed trails in $K'_{5,5}$ and $k \in [1, 2]$, then $|(V(T_1) \cup V(T_2)) \cap X_{1,5}^k| \geq 3$.*

Proof. If $|E(T_1) \cup E(T_2)| \geq 10$, then the edges of $E(T_1) \cup E(T_2)$ must cover at least $\lceil \frac{10}{4} \rceil = 3$ vertices of $X_{1,5}^k$ (note that $\Delta(K'_{5,5}) = 4$). The same is true if both T_1 and T_2 are 4-trails, since the subgraph of $K'_{5,5}$ that is induced by the eight edges incident with x_i^k or x_j^k , $i, j \in [1, 5]$, $i \neq j$, has two vertices of degree 1 (namely x_i^{3-k} and x_j^{3-k}), and so it cannot be equal to $T_1 \cup T_2$. ■

Theorem 9 *The graph G_a is ADCT for any odd integer $a \geq 7$. Moreover, given $s \in [4, 5]$, any sequence $\tau = (t_1, \dots, t_p) \in \text{Sct}(G_a)$ of length $p \geq 2$ has a G_a -realization (T_1, \dots, T_p) such that T_1 contains as a subgraph a 3-vertex path with endvertices x_1^2 and x_s^2 and T_2 contains the vertex x_2^2 .*

Proof. We use the general strategy with ADCT graphs $G^1 := K'_{5,5}$ (Proposition 5) and $G^2 := G_a^1$ (Theorem 3) the structure of the graph G_a is presented in Figure 2.




Figure 2: The graph G_a

First we show how to proceed provided three special conditions are fulfilled.

(C1) If there is I^1 with $[1, 2] \subseteq I^1 \subseteq [1, p]$ and $\sum_{i \in I^1} t_i = |E(G^1)| = 20$, put $I^2 := [1, p] - I^1$ and $\tau^l := \tau \setminus I^l$, $l = 1, 2$. There is a G^1 -realization $(T_1, T_2)\mathcal{T}^1$ of the sequence τ^1 and a G^2 -realization \mathcal{T}^2 of the sequence τ^2 . Then $\mathcal{T} := (T_1, T_2)\mathcal{T}^1\mathcal{T}^2$ is a G_a -realization of the sequence $\tau^1\tau^2 \sim \tau$. Any closed trail in a bipartite graph with bipartition $\{U, V\}$ is an alternating sequence of vertices of U and

V. Therefore, by Proposition 7 and Lemma 8, we may suppose without loss of generality that the trails T_1 and T_2 have the required properties.

(C2) If there are I^1 and $j \in [1, p] - I^1$ such that $[1, 2] \subseteq I^1 \cup \{j\}$, $\sum_{i \in I^1} t_i \leq 16$ and $\sum_{i \in I^1} t_i + t_j \geq 24$, put $I^2 := [1, p] - I^1 - \{j\}$, $t_j^1 := 20 - \sum_{i \in I^1} t_i$ and $t_j^2 := \sum_{i \in I^1} t_i + t_j - 20$. There is a G^l -realization $(T_j^l)\mathcal{T}^l$ of the sequence $(t_j^l)\tau\langle I^l \rangle \in \text{Sct}(G^l)$, $l = 1, 2$; for $i \in [1, 2] - \{j\} \subseteq I^1$ let T_i be a t_i -trail of \mathcal{T}^1 . Using Propositions 6, 7 and Lemma 8 we may suppose without loss of generality that T_1 (or T_1^1 if $j = 1$) contains as a subgraph a 3-vertex path with endvertices x_1^2 and x_s^2 , T_2 (or T_2^1 if $j = 2$) contains the vertex x_2^2 and $V(T_j^1) \cap V(T_j^2) \cap X_{1,4}^1 \neq \emptyset$. Then $T_j := T_j^1 \cup T_j^2$ is a t_j -trail and $(T_j)\mathcal{T}^1\mathcal{T}^2$ is an appropriate G_a -realization of the sequence $(t_j)\tau\langle I^1 \rangle\tau\langle I^2 \rangle \sim \tau$.

(C3) If there are I^1 and $\{j, k\} \subseteq [1, p] - I^1$ such that $[1, 2] \subseteq I^1 \cup \{j, k\}$, $\min\{t_j, t_k\} \geq 8$, $\sum_{i \in I^1} t_i \leq 12$ and $\sum_{i \in I^1} t_i + t_j + t_k \geq 28$, put $I^2 := [1, p] - I^1 - \{j, k\}$, $t_j^1 := \min\{16 - \sum_{i \in I^1} t_i, t_j - 4\}$, $t_k^1 := \max\{4, 24 - \sum_{i \in I^1} t_i - t_j\}$, $t_j^2 := t_j - t_j^1$ and $t_k^2 := t_k - t_k^1$. Then $t_j^1 + t_k^1 + \sum_{i \in I^1} t_i = |E(G^l)|$ and there is a G^l -realization $(T_j^l, T_k^l)\mathcal{T}^l$ of the sequence $(t_j^l, t_k^l)\tau\langle I^l \rangle$, $l = 1, 2$; for $i \in [1, 2] - \{j, k\} \subseteq I^1$ let T_i be a t_i -trail of \mathcal{T}^1 . By Propositions 6, 7 and Lemma 8 we may suppose without loss of generality that T_1 (or T_1^1 if $1 \in \{j, k\}$) contains as a subgraph a 3-vertex path with endvertices x_1^2 and x_s^2 , T_2 (or T_2^1 if $2 \in \{j, k\}$) contains the vertex x_2^2 and $V(T_m^1) \cap V(T_m^2) \cap X_{1,4}^1 \neq \emptyset$ for any $m \in \{j, k\}$. Then $T_m := T_m^1 \cup T_m^2$ is a t_m -trail, $m = j, k$ and $(T_j, T_k)\mathcal{T}^1\mathcal{T}^2$ is a required G_a -realization of the sequence $(t_j, t_k)\tau\langle I^1 \rangle\tau\langle I^2 \rangle \sim \tau$.

Let $i_1, i_2 \in [1, 2]$ be such that $i_1 \neq i_2$ and $t_{i_1} \leq t_{i_2}$. Since there are no additional requirements on T_i with $i \in [3, p]$, having in mind Lemma 1, in our analysis we may suppose without loss of generality that $t_i \leq t_{i+1}$ for any $i \in [3, p-1]$.

(1) $t_1 + t_2 \geq 24$

(11) If $t_{i_1} \geq 18$, then $I^1 := \emptyset$, $j := 1$, $k := 2 \rightarrow$ (C3), i.e. the condition (C3) is satisfied with the presented values of I^1 , j and k .

(12) If $t_{i_1} \leq 16$, then $I^1 := \{i_1\}$, $j := i_2 \rightarrow$ (C2).

(2) If $t_1 + t_2 = 22$, then $t_{i_1} \leq 10$, $t_{i_2} \geq 12$ and $\sum_{i=3}^p t_i = 4a - 22 \equiv 2 \pmod{4}$, hence there is $l \in [3, p]$ with $t_l \equiv 2 \pmod{4}$.

(21) If $t_p \geq 8$, then $I^1 := \{i_1\}$, $j := i_2$, $k := p \rightarrow$ (C3).

(22) If $t_p (= t_l) = 6$, then $I^1 := \{i_1, p\}$, $j := i_2 \rightarrow$ (C2).

(3) If $t_1 + t_2 = 20$, then $I^1 := [1, 2] \rightarrow$ (C1).

(4) If $t_1 + t_2 = 18$, then $t_{i_1} \leq 8$, $t_{i_2} \geq 10$ and there is $l \in [3, p]$ with $t_l \equiv 2 \pmod{4}$.

(41) If $t_l \geq 10$, then $I^1 := \{i_1\}$, $j := i_2$, $k := l \rightarrow$ (C3).

(42) If $t_l = 6$, then $I^1 := \{i_1, l\}$, $j := i_2 \rightarrow$ (C2).

(5) If $t_1 + t_2 \leq 16$, let $q \in [2, p-1]$ be determined by the inequalities $\sum_{i=1}^q t_i \leq 22$ and $\sum_{i=1}^{q+1} t_i \geq 24$.

(51) If $\sum_{i=1}^q t_i = 22$, then $q \geq 3$ and there is $l \in [q+1, p]$ with $t_l \equiv 2 \pmod{4}$.

(511) $t_q \geq 6$

(5111) If $t_p \geq t_q + 2$, then $I^1 := [1, q - 1]$, $j := p \rightarrow$ (C2).

(5112) If $t_i = t_q$ for any $i \in [q + 1, p]$, then $t_q = t_l \equiv 2 \pmod{4}$.

(51121) If $t_q \geq 10$, then $I^1 := [1, q - 1]$, $j := q$, $k := q + 1 \rightarrow$ (C3).

(51122) If $t_q = 6$, put $\tau^1 := (4) \prod_{i=1}^{q-1} (t_i) \in \text{Sct}(G^1)$, $\tau^2 := (8)(6)^{p-1-q} \in \text{Sct}(G^2)$ and consider a G^1 -realization $(T_q^1) \prod_{i=1}^{q-1} (T_i)$ of the sequence τ^1 and a G^2 -realization $(T_{q+1}^2) \prod_{i=q+2}^p (T_i)$ of the sequence τ^2 yielded by Theorem 4. Let $T_q^1 = \prod_{i=1}^5 (b_i)$ with $b_1 = b_5 \in X_{1,5}^1$ and let $T_{q+1}^2 = \prod_{i=1}^9 (c_i)$ with $c_1 = c_9 \in X_{1,4}^1$. Since T_{q+1}^2 is a cycle, we have $V(T_{q+1}^2) \cap X_{1,4}^1 = X_{1,4}^1$. By Proposition 7 we may suppose without loss of generality that $b_1 = c_1$ and $b_3 = c_5$. With $T_q := (c_1, b_2) \prod_{i=5}^9 (c_i)$ and $T_{q+1} := (c_1, b_4) \prod_{i=1}^5 (c_{6-i})$ then (T_1, \dots, T_p) is a G_a -realization of the sequence τ . Since $q \geq 3$, by Proposition 7 and Lemma 8 we may suppose without loss of generality that the additional requirements on T_1 and T_2 are fulfilled.

(512) If $t_q = 4$, then $t_1 + t_2 \equiv 2 \pmod{4}$, and so $q \geq 4$ and $\sum_{i=1}^{q-2} t_i = 14$.

(5121) If $t_p \geq 10$, then $I^1 := [1, q - 2]$, $j := p \rightarrow$ (C2).

(5122) If $t_p \leq 8$, then $t_l = 6$ and $I^1 := [1, q - 2] \cup \{l\} \rightarrow$ (C1).

(52) If $\sum_{i=1}^q t_i = 20$, then $I^1 := [1, q] \rightarrow$ (C1).

(53) If $\sum_{i=1}^q t_i = 18$, then $q \geq 3$ and there is $l \in [q + 1, p]$ with $t_l \equiv 2 \pmod{4}$.

(531) If $t_q \geq 6$, then $\sum_{i=1}^{q-1} t_i \leq 12$.

(5311) If $t_p \geq t_q + 6$, then $I^1 := [1, q - 1]$, $j := p \rightarrow$ (C2).

(5312) If there is $m \in [q + 1, p]$ with $t_m = t_q + 2$, then $I^1 := [1, q - 1] \cup \{m\} \rightarrow$ (C1).

(5313) If $t_i \in \{t_q, t_q + 4\}$ for any $i \in [q + 1, p]$, then $t_q \equiv t_l \equiv 2 \pmod{4}$, hence $t_q \leq 10$.

(53131) If $t_q = 10$, then $q = 3$, $I^1 := [1, q - 1]$, $j := q$, $k := q + 1 \rightarrow$ (C3).

(53132) If $t_q = 6$, put $\tau^1 := (8) \prod_{i=1}^{q-1} (t_i) \in \text{Sct}(G^1)$ and $\tau^2 := (t_p - 2) \prod_{i=q+1}^{p-1} (t_i) \in \text{Sct}(G^2)$. Consider a G^1 -realization $(T_q^1) \prod_{i=1}^{q-1} (T_i)$ of the sequence τ^1 and a G^2 -realization $(T_p^2) \prod_{i=q+1}^{p-1} (T_i)$ of the sequence τ^2 . Let $T_q^1 = \prod_{i=1}^9 (b_i)$ with $b_1 = b_9 \in X_{1,5}^1$ and let $T_p^2 = \prod_{i=1}^{t_p-1} (c_i)$ with $c_1 = c_{t_p-1} \in X_{1,4}^1$. We have $|V(T_q^1) \cap X_{1,5}^1| \geq 3$ (if T_q^1 is not a cycle, it is a union of two edge-disjoint 4-trails and then it suffices to use Lemma 8). Therefore, we may suppose without loss of generality that $b_5 \neq b_1$. Moreover, by Proposition 6, the assumption $c_1 = b_1$ and $c_3 = b_5$ also does not cause a loss of generality. With $T_q := (b_1, c_2) \prod_{i=1}^5 (b_{6-i})$ and $T_p := (c_1, b_8, b_7, b_6) \prod_{i=3}^{t_p-1} (c_i)$ then, using Proposition 7 and Lemma 8, we may suppose without loss of generality that (T_1, \dots, T_p) is an appropriate G_a -realization of the sequence τ .

(532) $t_q = 4$

(5321) If $t_l \geq 10$, then $I^1 := [1, q - 1]$, $j := l \rightarrow$ (C2).

(5322) If $t_l = 6$, then $I^1 := [1, q - 1] \cup \{l\} \rightarrow$ (C1).

(54) If $\sum_{i=1}^q t_i \leq 16$, then $I^1 := [1, q]$, $j := q + 1 \rightarrow$ (C2). ■

Theorem 10 *The graph H_a is ADCT for any odd integer $a \geq 7$. Moreover, any sequence $\tau = (t_1, \dots, t_p) \in \text{Sct}(H_a)$ of length $p \geq 2$ has an H_a -realization (T_1, \dots, T_p) such that there are $(i_r, j_r) \in [5, a] \times [1, 2]$ with $x_{i_r}^{j_r} \in V(T_r)$, $r = 1, 2$, and $i_1 \neq i_2$.*

Proof. We proceed similarly as in the proof of Theorem 9 with ADCT graphs $G^1 := G_a^2$ (Theorem 3) and $G^2 := G_a$ (Theorem 9). The graph H_a is depicted in Figure 3.




Figure 3: The graph H_a

(C4) If there is $I^1 \subseteq [1, p]$ such that $|[1, 2] \cap I^1| \geq 1$ and $\sum_{i \in I^1} t_i = |E(G^1)| = 4a - 20$, put $I^2 := [1, p] - I^1$ and $\tau^l := \tau \langle I^l \rangle$, $l = 1, 2$. Let \mathcal{T}^l be a G^l -realization of the sequence τ^l , $l = 1, 2$, and let T_i be a t_i -trail of $\mathcal{T}^1 \mathcal{T}^2$, $i = 1, 2$. If $[1, 2] \subseteq I^1$, by Proposition 6 we may suppose without loss of generality that $x_{5+i}^1 \in V(T_i)$, $i = 1, 2$; in such a case we are done with $(i_1, j_1) := (6, 1)$ and $(i_2, j_2) := (7, 1)$. If there is $m \in [1, 2]$ such that $m \in I^1$ and $3 - m \in I^2$, then, by Proposition 6 and Theorem 9, we may suppose without loss of generality that $(i_m, j_m) := (6, 1)$ and $(i_{3-m}, j_{3-m}) := (5, 2)$ are appropriate pairs.

(C5) If there are I^1 and $j \in [1, p] - I^1$ such that $|[1, 2] \cap (I^1 \cup \{j\})| \geq 1$, $\sum_{i \in I^1} t_i \leq 4a - 24$ and $\sum_{i \in I^1} t_i + t_j \geq 4a - 16$, put $I^2 := [1, p] - I^1 - \{j\}$, $t_j^1 := 4a - 20 - \sum_{i \in I^1} t_i$, $t_j^2 := \sum_{i \in I^1} t_i + t_j + 20 - 4a$ and $m := \min(\{0\} \cup I^2)$. Consider a G^1 -realization $(T_j^1) \mathcal{T}^1$ of the sequence $(t_j^1) \tau \langle I^1 \rangle \in \text{Sct}(G^1)$ and let T_i be a t_i -trail of \mathcal{T}^1 with $i \in ([1, 2] - \{j\}) \cap I^1$. By Proposition 6 we may suppose without loss of generality that $x_2^2 \in V(T_j^1)$, $j \in [1, 2] \Rightarrow x_{5+j}^1 \in V(T_j^1)$ and $x_{5+i}^1 \in V(T_i)$ for any $i \in ([1, 2] - \{j\}) \cap I^1$.

If $I^2 \neq \emptyset$ (so that $m \geq 1$), by Theorem 9 there is a G^2 -realization $(T_m, T_j^2) \mathcal{T}_2$ of the sequence $(t_m, t_j^2) \tau \langle I^2 - \{m\} \rangle \in \text{Sct}(G^2)$ such that $\{x_1^2, x_5^2\} \subseteq V(T_m)$ and

$x_2^2 \in V(T_j^2)$. Then $T_j := T_j^1 \cup T_j^2$ is a t_j -trail and $(T_j, T_m)\mathcal{T}^1\mathcal{T}^2$ is a required H_a -realization of the sequence $(t_j, t_m)\tau\langle I^1 \rangle\tau\langle I^2 - \{m\} \rangle \sim \tau$. Appropriate pairs are as follows: if $m \in [1, 2]$, then $(i_m, j_m) := (5, 2)$ and $(i_{3-m}, j_{3-m}) := (8 - m, 1)$; if $m \notin [1, 2]$, then $(i_r, j_r) := (5 + r, 1)$, $r = 1, 2$.

If $I^2 = \emptyset$ (and $m = 0$), then $T_j := T_j^1 \cup G^2$ is a t_j -trail and $(T_j^1)\mathcal{T}_1$ is an appropriate H_a -realization of the sequence $(t_j)\tau\langle I^1 \rangle \sim \tau$.

(C6) If there are I^1 and $\{j, k\} \subseteq [1, p] - I^1$ such that $[1, 2] \subseteq I^1 \cup \{j, k\}$, $\min\{t_j, t_k\} \geq 8$, $\sum_{i \in I^1} t_i \leq 4a - 28$ and $\sum_{i \in I^1} t_i + t_j + t_k \geq 4a - 12$ (we may suppose without loss of generality that $j < k$), then with $I^2 := [1, p] - I^1 - \{j, k\}$, $t_j^1 := \min\{4a - 24 - \sum_{i \in I^1} t_i, t_j - 4\}$, $t_k^1 := \max\{4, 4a - 16 - \sum_{i \in I^1} t_i - t_j\}$, $t_j^2 := t_j - t_j^1$ and $t_k^2 := t_k - t_k^1$ we have $t_j^1 + t_k^1 + \sum_{i \in I^1} t_i = |E(G^l)|$ and $\tau^l := (t_j^l, t_k^l)\tau\langle I^l \rangle \in \text{Sct}(G^l)$, $l = 1, 2$. Consider a G^1 -realization $(T_j^1, T_k^1)\mathcal{T}^1$ of the sequence τ^1 and let T_i be a t_i -trail of \mathcal{T}^1 with $i \in [1, 2] - \{j, k\} \subseteq I^1$. Because of Proposition 6 we may suppose without loss of generality that $x_1^2 \in V(T_j^1)$, $x_2^2 \in V(T_k^1)$, $m \in [1, 2] \cap \{j, k\} \Rightarrow x_{5+m}^1 \in V(T_m^1)$ and $x_{5+i}^1 \in V(T_i)$ for any $i \in [1, 2] - \{j, k\}$. By Theorem 9 there is a G^2 -realization $(T_j^2, T_k^2)\mathcal{T}^2$ of the sequence τ^2 such that $x_1^2 \in V(T_j^2)$ and $x_2^2 \in V(T_k^2)$. Then $T_m := T_m^1 \cup T_m^2$ is a t_m -trail, $m = j, k$ and $(T_j, T_k)\mathcal{T}^1\mathcal{T}^2$ is an H_a -realization of the sequence $(t_j, t_k)\tau\langle I^1 \rangle\tau\langle I^2 \rangle \sim \tau$ with required properties; appropriate pairs are $(i_r, j_r) := (5 + r, 1)$, $r = 1, 2$.

The additional requirements on T_1 and T_2 are symmetrical and there are no additional requirements on T_i with $i \in [3, p]$; therefore, in our analysis we may suppose without loss of generality that $t_1 \leq t_2$ and $t_i \leq t_{i+1}$ for any $i \in [3, p - 1]$.

(1) $t_1 + t_2 \geq 4a - 16$

(11) If $t_1 \leq 4a - 24$, then $I^1 := \{1\}$, $j := 2 \rightarrow$ (C5).

(12) If $t_1 \geq 4a - 22$, then $t_1 \geq 6$.

(121) If $a \geq 9$, then $t_1 + t_2 \geq 8a - 44 \geq 4a - 12$, $t_1 \geq 14$ and $I^1 := \emptyset$, $j := 1$, $k := 2 \rightarrow$ (C6).

(122) If $a = 7$, then $|E(G^1)| = 8$.

(1221) If $t_1 \geq 8$, then $t_1 + t_2 \geq 4a - 12$ and $I^1 := \emptyset$, $j := 1$, $k := 2 \rightarrow$ (C6).

(1222) If $t_1 = 6$, by Theorem 9 there is a G^2 -realization $(T_2^2) \prod_{i=3}^p (T_i)$ of the sequence $(t_2 - 2) \prod_{i=3}^p (t_i) \in \text{Sct}(G^2)$ such that T_2^2 contains as a subgraph a 3-vertex path with endvertices x_1^2 and x_4^2 . Thus, we may suppose without loss of generality that $T_2^2 = \prod_{i=1}^{t_2-1} (c_i)$ where $c_1 = c_{t_2-1} = x_1^2$ and $c_3 = x_4^2$. With $T_1 := (x_1^2, c_2, x_4^2, x_7^1, x_3^2, x_6^1, x_1^2)$ and $T_2 := (c_1, x_7^1, x_2^2, x_6^1) \prod_{i=3}^{t_2-1} (c_i)$ then (T_1, \dots, T_p) is a required H_a -realization of the sequence τ ; appropriate pairs are $(i_r, j_r) := (5 + r, 1)$, $r = 1, 2$.

(2) If $t_1 + t_2 = 4a - 18$, then $\sum_{i=3}^p t_i = 4a - 2 \equiv 2 \pmod{4}$ and there is $l \in [3, p]$ satisfying $t_l \equiv 2 \pmod{4}$.

(21) If $t_1 \leq 4a - 28$, then $t_2 \geq 10$.

(211) If $t_p \geq 8$, then $I^1 := \{1\}$, $j := 2$, $k := p \rightarrow$ (C6).

(212) $t_p (= t_l) = 6$

(2121) If $t_1 \leq 4a - 30$, then $I^1 := \{1, p\}$, $j := 2 \rightarrow$ (C5).

(2122) If $t_1 = 4a - 28$, then $t_2 = 10$, $a \leq 9$, $t_1 = 8$, $a = 9$ and $I^1 := \{2, p\} \rightarrow$ (C4).

(3) If $t_1 + t_2 = 4a - 20$, then $I^1 := [1, 2] \rightarrow$ (C4).

(4) If $t_1 + t_2 = 4a - 22$, then $a \geq 9$, $t_2 \geq 8$ and there is $l \in [3, p]$ with $t_l \equiv 2 \pmod{4}$.

(41) If $t_1 \leq 4a - 34$, then $t_2 \geq 12$.

(411) If $t_l \geq 10$, then $I^1 := \{1\}$, $j := 2$, $k := l \rightarrow$ (C6).

(412) If $t_l = 6$, then $I^1 := \{1, l\}$, $j := 2 \rightarrow$ (C5).

(42) If $t_1 \geq 4a - 32$, then $a = 9$ and $t_2 \in \{8, 10\}$.

(421) If $t_l \geq 10$, then $I^1 := \{1\}$, $j := 2$, $k := l \rightarrow$ (C6).

(422) If $t_l = 6$, then $t_i \in \{4, 6\}$ for any $i \in [3, p]$, $\sum_{i=3}^p t_i = 38$ and the sequence $\prod_{i=3}^p (t_i)$ contains at least two 4's and at least one 6. Thus, there is $I^1 \subseteq [2, p]$ such that $2 \in I^1$, $\sum_{i \in I^1} t_i = 16$ and the condition (C4) is satisfied.

(5) If $t_1 + t_2 \leq 4a - 24$, let $q \in [2, p - 1]$ be determined by the inequalities $\sum_{i=1}^q t_i \leq 4a - 18$ and $\sum_{i=1}^{q+1} t_i \geq 4a - 16$.

(51) If $\sum_{i=1}^q t_i = 4a - 18$, then $q \geq 3$ and there is $l \in [q + 1, p]$ with $t_l \equiv 2 \pmod{4}$.

(511) $t_q \geq 6$

(5111) If $t_p \geq t_q + 2$, then $I^1 := [1, q - 1]$, $j := p \rightarrow$ (C5).

(5112) If $t_i = t_q$ for any $i \in [q + 1, p]$, then $t_q = t_l \equiv 2 \pmod{4}$.

(51121) If $t_q \geq 10$, then $I^1 := [1, q - 1]$, $j := q$, $k := q + 1 \rightarrow$ (C6).

(51122) If $t_q = 6$, then $6|4a - 2 = 6(p - q)$, hence $a \equiv 5 \pmod{6}$ and $p - q \geq 7$.

(511221) If $t_2 \geq 12$, then $I^1 := \{1\} \cup [3, q + 1]$, $j := 2 \rightarrow$ (C5).

(511222) $t_2 \leq 10$

(5112221) If $t_2 = 10$, then $I^1 := [q + 5, p]$, $j := 2 \rightarrow$ (C5).

(5112222) If $t_2 = 8$, then $I^1 := \{1\} \cup [3, q + 1] \rightarrow$ (C4).

(5112223) If $t_2 = 6$, then $I^1 := \{2\} \cup [q + 5, p] \rightarrow$ (C4).

(5112224) $t_2 = 4$

(51122241) If $t_3 = 4$, then $I^1 := [1, 3] \cup [q + 6, p] \rightarrow$ (C4).

(51122242) If $t_3 = 6$, then $\tau = (4)^2(6)^{p-2}$, $6p - 4 = |E(H_a)| = 8a - 20$ and $p \equiv 0 \pmod{2}$. Put $\tau_1 := (8)(6)^2$, $\tau_2 := (6)^{\frac{p-4}{2}} =: \tau_3$ and consider a $K_{5,5}^1$ -realization $(T_{1,2}, T_3, T_4)$ of the sequence τ_1 presented in Figure 1, a G_a^1 -realization $(\tilde{T}_5) \prod_{i=6}^{\frac{p+4}{2}} (T_i)$ of the sequence τ_2 and a G_a^2 -realization $\prod_{i=\frac{p+6}{2}}^p (T_i)$ of the sequence τ_3 . The closed trail $T_{1,2}$ is an 8-cycle, hence by Proposition 7 we may suppose without loss of generality that $V(T_{1,2}) \cap X_{1,5}^1 = X_{1,4}^1$ and $T_{1,2} = \prod_{i=1}^9 (b_i)$ with $b_1 = b_9 \in X_{1,4}^1$. By Proposition 6 we may suppose without loss of generality that $\tilde{T}_5 = \prod_{i=1}^7 (c_i)$ with $c_1 = c_7 = b_1$, $c_3 = b_3$, $c_5 = b_7$, $c_2 = x_6^2$ and $c_6 = x_7^2$. Then (T_1, \dots, T_p) with $T_1 := (b_1, b_2, b_3, c_2, b_1)$, $T_2 := (b_9, b_8, b_7, c_6, b_9)$ and $T_5 := (b_3, c_4, b_7, b_6, b_5, b_4, b_3)$ is a required H_a -realization of the sequence τ ; appropriate pairs are $(i_r, j_r) := (5 + r, 2)$, $r = 1, 2$.

(512) If $t_q = 4$, then $q \geq 4$ and $\sum_{i=1}^{q-2} t_i = 4a - 26$.

(5121) If $t_p \geq 10$, then $I^1 := [1, q - 2]$, $j := p \rightarrow$ (C5).

- (5122) If $t_p \leq 8$, then $t_l = 6$ and $I^1 := [1, q - 2] \cup \{l\} \rightarrow (C4)$.
- (52) If $\sum_{i=1}^q t_i = 4a - 20$, then $I^1 := [1, q] \rightarrow (C4)$.
- (53) If $\sum_{i=1}^q t_i = 4a - 22$, then $q \geq 3$.
- (531) $t_q \geq 6$
- (5311) If $t_p \geq t_q + 6$, then $I^1 := [1, q - 1]$, $j := p \rightarrow (C5)$.
- (5312) If there is $m \in [q+1, p]$ such that $t_m = t_q + 2$, then $I^1 := [1, q - 1] \cup \{m\} \rightarrow (C4)$.
- (5313) If $t_i \in \{t_q, t_q + 4\}$ for any $i \in [q + 1, p]$, then $t_q \equiv t_l \equiv 2 \pmod{4}$, $(p - q)(t_q + 4) \geq 4a + 2 = \sum_{i=1}^q t_i + 24 \geq t_q + 24$, $p - q \geq \frac{t_q + 24}{t_q + 4} > 1$ and $p - q \geq 2$.
- (53131) If $t_{p-1} \geq 10$, then $I^1 := [1, q - 1]$, $j := p - 1$, $k := p \rightarrow (C6)$.
- (53132) If $t_{p-1} = 6$, then $t_q = 6$.
- (531321) If $t_2 \geq 8$, then $I^1 := \{1\} \cup [3, q + 1]$, $j := 2 \rightarrow (C5)$.
- (531322) If $t_2 \leq 6$, then by Theorem 4 there exists a G^1 -realization $\mathcal{T}^1 := (T_q^1) \prod_{i=1}^{q-1} (T_i)$ of the sequence (8) $\prod_{i=1}^{q-1} (t_i)$ such that all trails of \mathcal{T}^1 are cycles. Therefore, by Proposition 6 we may suppose without loss of generality that $x_{5+i}^1 \in V(T_i)$, $i = 1, 2$, and $T_q^1 = \prod_{i=1}^9 (b_i)$ with $b_1 = b_9 = x_1^2$ and $b_5 = x_4^2$. By Theorem 9 there is a G^2 -realization $(T_{q+1}^2) \prod_{i=q+2}^p (T_i)$ of the sequence (4) $\prod_{i=q+2}^p (t_i)$ such that T_{q+1}^2 contains as a subgraph a 3-vertex path with endvertices x_1^2 and x_4^2 . Thus, we may suppose without loss of generality that $T_{q+1}^2 = \prod_{i=1}^5 (c_i)$ where $c_1 = c_5 = x_1^2$ and $c_3 = x_4^2$. Then (T_1, \dots, T_p) with $T_{q+1} := (b_5, c_4) \prod_{i=1}^5 (b_i)$ and $T_{q+2} := (b_9, c_2) \prod_{i=5}^9 (b_i)$ is a required H_a -realization of the sequence τ ; appropriate pairs are $(i_r, j_r) := (5 + r, 1)$, $r = 1, 2$.
- (532) $t_q = 4$
- (5321) If $t_p \geq 10$, then $I^1 := [1, q - 1]$, $j := p \rightarrow (C5)$.
- (5322) If $t_p \leq 8$, then $t_l = 6$ and $I^1 := [1, q - 1] \cup \{l\} \rightarrow (C4)$.
- (54) If $\sum_{i=1}^q t_i \leq 4a - 24$, then $I^1 := [1, q]$, $j := q + 1 \rightarrow (C5)$. ■

Theorem 11 *If a is an odd integer, $a \geq 3$, then the graph $K'_{a,a}$ is ADCT. Moreover, if $r = \frac{a(a-1)-2}{6} \in \mathbb{Z}$, there is a $K'_{a,a}$ -realization (T_1, \dots, T_r) of the sequence $(6)^{r-1}(8) \in \text{Sct}(K'_{a,a})$ such that T_r has as a subgraph a 5-vertex path.*

Proof. We proceed by induction on a . The graphs $K'_{a,a}$ with $a \leq 5$ are ADCT by Proposition 5. Further, the 8-trail of the $K'_{5,5}$ -realization of the sequence $(6)^2(8) \in \text{Sct}(K'_{5,5})$ presented in Figure 1 is a cycle, and so trivially it has as a subgraph a 5-vertex path.

So, suppose that $a \geq 7$, the graph $K'_{a-4,a-4}$ is ADCT and, provided $s := \frac{(a-4)(a-5)-2}{6} \in \mathbb{Z}$, there is a G^1 -realization $\prod_{i=1}^s (T_i^1)$ of the sequence $(6)^{s-1}(8) \in \text{Sct}(G^1)$ such that T_s^1 has as a subgraph a 5-vertex path. We can use again the general strategy, since the graph $K'_{a,a}$ (see Figure 4) is an edge-disjoint union of ADCT graphs $G^1 := F_a$ (the induction hypothesis) and $G^2 := H_a$ (Theorem 10). Consider a sequence $\tau = (t_1, \dots, t_p) \in \text{Sct}(K'_{a,a})$.




Figure 4: The graph $K'_{a,a}$

(C7) If there is $I^1 \subseteq [1, p]$ such that $\sum_{i \in I^1} t_i = a^2 - 9a + 20 = |E(G^1)|$, put $I^2 := [1, p] - I^1$, $\tau^l := \tau \langle I^l \rangle \in \text{Sct}(G^l)$ and consider a G^l -realization \mathcal{T}^l of the sequence τ^l , $l = 1, 2$. Then $\mathcal{T}^1 \mathcal{T}^2$ is a $K'_{a,a}$ -realization of the sequence $\tau^1 \tau^2 \sim \tau$.

(C8) If there are I^1 and $j \in [1, p] - I^1$ such that $\sum_{i \in I^1} t_i \leq a^2 - 9a + 16$ and $\sum_{i \in I^1} t_i + t_j \geq a^2 - 9a + 24$, put $I^2 := [1, p] - I^1 - \{j\}$, $t_j^1 := a^2 - 9a + 20 - \sum_{i \in I^1} t_i$, $t_j^2 := \sum_{i \in I^1} t_i + t_j - a^2 + 9a - 20$. Then $\tau^l := (t_j^l) \tau \langle I^l \rangle \in \text{Sct}(G^l)$, $l = 1, 2$. By Theorem 10 there is a G^2 -realization $(T_j^2) \mathcal{T}^2$ of the sequence τ^2 such that there is $(i_1, j_1) \in [5, a] \times [1, 2]$ with $x_{i_1}^{j_1} \in V(T_j^2)$. By the induction hypothesis there is a G^1 -realization $(T_j^1) \mathcal{T}^1$ of the sequence τ^1 ; by Proposition 7 we may suppose without loss of generality that $x_{i_1}^{j_1} \in V(T_j^1)$. Then $T_j := T_j^1 \cup T_j^2$ is a t_j -trail and $(T_j) \mathcal{T}^1 \mathcal{T}^2$ is a $K'_{a,a}$ -realization of the sequence $(t_j) \tau \langle I^1 \rangle \tau \langle I^2 \rangle \sim \tau$.

(C9) If there are I^1 and $\{j, k\} \subseteq [1, p] - I^1$ such that $\min\{t_j, t_k\} \geq 8$, $\sum_{i \in I^1} t_i \leq a^2 - 9a + 12$ and $\sum_{i \in I^1} t_i + t_j + t_k \geq a^2 - 9a + 28$, then with $I^2 := [1, p] - I^1 - \{j, k\}$, $t_j^1 := \min\{a^2 - 9a + 16 - \sum_{i \in I^1} t_i, t_j - 4\}$, $t_k^1 := \max\{4, a^2 - 9a + 24 - \sum_{i \in I^1} t_i - t_j\}$, $t_j^2 := t_j - t_j^1$ and $t_k^2 := t_k - t_k^1$ we have $t_j^1 + t_k^1 + \sum_{i \in I^1} t_i = |E(G^l)|$ and $\tau^l := (t_j^l, t_k^l) \tau \langle I^l \rangle \in \text{Sct}(G^l)$, $l = 1, 2$. Theorem 10 yields a G^2 -realization $(T_j^2, T_k^2) \mathcal{T}^2$ of the sequence τ^2 such that there are $(i_r, j_r) \in [5, a] \times [1, 2]$, $r = 1, 2$, with $x_{i_1}^{j_1} \in V(T_j^2)$, $x_{i_2}^{j_2} \in V(T_k^2)$ and $i_1 \neq i_2$. By the induction hypothesis there is a G^1 -realization $(T_j^1, T_k^1) \mathcal{T}^1$ of the sequence τ^1 ; by Proposition 7 we may suppose without loss of generality that $x_{i_1}^{j_1} \in V(T_j^1)$ and $x_{i_2}^{j_2} \in V(T_k^1)$ (note that both T_j^1 and T_k^1 have at least two vertices in both $X_{5,a}^1$ and $X_{5,a}^2$). Then $T_m := T_m^1 \cup T_m^2$ is a t_m -trail, $m = j, k$ and $(T_j, T_k) \mathcal{T}^1 \mathcal{T}^2$ is a $K'_{a,a}$ -realization of the sequence $(t_j, t_k) \tau \langle I^1 \rangle \tau \langle I^2 \rangle \sim \tau$.

Because of Lemma 1 we may suppose without loss of generality that τ is a

nondecreasing sequence. Let $q \in [0, p - 1]$ be determined by the inequalities $\sum_{i=1}^q t_i \leq a^2 - 9a + 22$ and $\sum_{i=1}^{q+1} t_i \geq a^2 - 9a + 24$.

(1) If $\sum_{i=1}^q t_i = a^2 - 9a + 22$, then $\sum_{i=q+1}^p t_i = 8a - 22$ and there is $l \in [q+1, p]$ such that $t_l \equiv 2 \pmod{4}$.

(11) $t_q \geq 6$

(111) If $t_p \geq t_q + 2$, then $I^1 := [1, q - 1]$, $j := p \rightarrow$ (C8).

(112) If $t_i = t_q$ for any $i \in [q + 1, p]$, then $t_q = t_l \equiv 2 \pmod{4}$.

(1121) If $t_q \geq 10$, then $I^1 := [1, q - 1]$, $j := q$, $k := q + 1 \rightarrow$ (C9).

(1122) If $t_q = 6$, then $6q \geq \sum_{i=1}^q t_i \geq 8$, $q \geq 2$, $8a - 22 = 6(p - q)$, $4a - 11 \equiv 0 \pmod{3}$, $a \equiv 5 \pmod{6}$, $a(a - 1) \equiv 2 \pmod{6}$, the sequence τ must contain at least two 4's and $I^1 := [3, q + 1] \rightarrow$ (C7).

(12) If $t_q = 4$, then $4q \geq 8$ and $q \geq 2$.

(121) If $t_l \geq 10$, then $I^1 := [1, q - 2]$, $j := l \rightarrow$ (C8).

(122) If $t_l = 6$, then $I^1 := [1, q - 2] \cup \{l\} \rightarrow$ (C7).

(2) If $\sum_{i=1}^q t_i = a^2 - 9a + 20$, then $I^1 := [1, q] \rightarrow$ (C7). Note that if r defined in the statement of our Theorem is integer, then $a(a - 1) \equiv 2 \pmod{6}$, $a \equiv 5 \pmod{6}$, $a^2 - 9a + 20 \equiv 0 \pmod{6}$, $4a - 20 \equiv 0 \pmod{6}$, and so $\tau = (6)^{p-1}(8)$ yields $8a - 20 = 6(p - q - 1) + 8$, $p - q - 1 \geq 60$, $6(p - q - 1) \equiv 0 \pmod{4}$ and $p - q - 1 \equiv 0 \pmod{2}$. The graph G^2 is an edge-disjoint union of ADCT graphs $G_1^2 := G_a^1$, $G_2^2 := G_a^2$ and $G_3^2 := K'_{5,5}$. Put $\tau^1 := (6)^q$, $\tau_1^2 := (6)^{\frac{p-q-3}{2}} =: \tau_2^2$, $\tau_3^2 := (6)^2(8)$ and let \mathcal{T}^1 be a G^1 -realization of the sequence τ^1 and let \mathcal{T}_m^2 be a G_m^2 -realization of the sequence τ_m^2 , $m = 1, 2, 3$, where $\mathcal{T}_3^2 = (T_{p-2}, T_{p-1}, T_p)$ is that presented in Figure 1. Then $\mathcal{T}^1 \mathcal{T}_1^2 \mathcal{T}_2^2 \mathcal{T}_3^2$ is a $K'_{a,a}$ -realization of the sequence $(6)^{p-1}(8)$ and the 8-trail T_p (that is a cycle) has trivially as a subgraph a 5-vertex path.

(3) If $\sum_{i=1}^q t_i = a^2 - 9a + 18$, there is $l \in [q + 1, p]$ such that $t_l \equiv 2 \pmod{4}$.

(31) $t_q \geq 6$

(311) If $t_p \geq t_q + 6$, then $I^1 := [1, q - 1]$, $j := p \rightarrow$ (C8).

(312) If there is $m \in [q + 1, p]$ such that $t_m = t_q + 2$, then $I^1 := [1, q - 1] \cup \{m\} \rightarrow$ (C7).

(313) If $t_i \in \{t_q, t_q + 4\}$ for any $i \in [q + 1, p]$, then $t_q \equiv t_l \equiv 2 \pmod{4}$.

(3131) $p \geq q + 2$

(31311) If $t_{p-1} \geq 10$, then $I^1 := [1, q - 1]$, $j := p - 1$, $k := p \rightarrow$ (C9).

(31312) $t_{p-1} = 6$

(313121) If $t_1 = 4$, then $I^1 := [2, q + 1] \rightarrow$ (C7).

(313122) If $t_1 = 6$, then $a^2 - 9a + 18 = 6q$, $a \equiv 3 \pmod{6}$, $\sum_{i=q+1}^p t_i = 8a - 18 \equiv 0 \pmod{6}$, $t_p = 6$, $\tau = (6)^p$, $8a - 18 = 6(p - q)$, $p - q \geq 9$, $6(p - q) \equiv 6 \pmod{48}$ and $p - q - 1 \equiv 0 \pmod{8}$. The graph G^2 is an edge-disjoint union of ADCT graphs $G_1^2 := G_a$ and $G_2^2 := G_a^2$. Put $\tau^1 := (8)(6)^{q-1}$, $\tau_1^2 := (6)^{\frac{p-q+3}{2}}$ and $\tau_2^2 := (4)(6)^{\frac{p-q-5}{2}}$. By the induction hypothesis and by Lemma 1 there is a G^1 -realization $(T_q^1) \mathcal{T}^1$ of the sequence τ^1 such that T_q^1 has as a subgraph a 5-vertex path. By Proposition 7 we may suppose without loss of generality that

$T_q^1 = \prod_{i=1}^9(b_i)$ where $b_1 = b_9 \in X_{5,a}^1$ and $\prod_{i=1}^5(b_i)$ is a path. By Theorem 10 there is a G_1^2 -realization \mathcal{T}_1^2 of the sequence τ_1^2 . Further, by Theorem 3 there is a G_2^2 -realization $(T_{q+1}^2)\mathcal{T}_2^2$ of the sequence τ_2^2 ; by Proposition 6 we may suppose without loss of generality that $T_{q+1}^2 = \prod_{i=1}^5(c_i)$ where $c_1 = c_5 = b_1$ and $c_3 = b_5$. With $T_q := (b_5, c_2) \prod_{i=1}^5(b_i)$ and $T_{q+1} := (b_9, c_4) \prod_{i=5}^9(b_i)$ then $(T_q, T_{q+1})\mathcal{T}^1\mathcal{T}_1^2\mathcal{T}_2^2$ is a $K'_{a,a}$ -realization of the sequence $\tau = (6)^p$.

(3132) If $p = q + 1$, then $t_p = 8a - 18$, $t_q \geq 8a - 22$ and $I^1 := [1, q - 1]$, $j := q$, $k := p \rightarrow$ (C9).

(32) $t_q = 4$

(321) If $t_l \geq 10$, then $I^1 := [1, q - 1]$, $j := l \rightarrow$ (C8).

(322) If $t_l = 6$, then $I^1 := [1, q - 1] \cup \{l\} \rightarrow$ (C7).

(4) If $\sum_{i=1}^q \leq a^2 - 9a + 16$, then $I^1 := [1, q]$, $j := q + 1 \rightarrow$ (C8). ■

References

- [1] P.N. Balister: Packing circuits into K_N . *Comb. Probab. Comput.* 6(2001), 463–499. Zbl pre01718972
- [2] P.N. Balister: Packing closed trails into dense graphs. *J. Comb. Theory, Ser. B* 88(2003), 107–118. Zbl 1045.05074
- [3] P.N. Balister: Packing digraphs with directed closed trails. *Comb. Probab. Comput.* 12(2003), 1–15. Zbl 1015.05072
- [4] Ch.-Ch. Chou, Ch.-M. Fu, W.-Ch. Huang: Decomposition of $K_{n,m}$ into short cycle. *Discrete Math.* 197/198(1999), 195–203. Zbl 0934.05106
- [5] S. Cichacz: Decomposition of complete bipartite digraphs and even complete bipartite multigraphs into closed trails. Preprint MD 016 (2005), <http://www.ii.uj.edu.pl/preMD/index.php>
- [6] S. Cichacz, J. Przybyło, M. Woźniak: Decompositions of pseudographs into closed trails of even sizes (submitted)
- [7] M. Horňák, Z. Kocková: On complete tripartite graphs arbitrarily decomposable into closed trails. *Tatra Mt. Math. Publ.* (to appear)
- [8] M. Horňák, M. Woźniak: Decomposition of complete bipartite even graphs into closed trails. *Czech. Math. J.* 53(128)(2003), 127–134. Zbl 1010.05054

Recent IM Preprints, series A

2003

- 1/2003 Ceclárová K.: *Eigenvectors of interval matrices over max-plus algebra*
2/2003 Mihók P. and Semanišin G.: *On invariants of hereditary graph properties*
3/2003 Ceclárová K.: *A problem on optimal transportation*

2004

- 1/2004 Jendroľ S. and Voss H.-J.: *Light subgraphs of graphs embedded in the plane and in the projective plane – survey*
2/2004 Dražnová S., Ivančo J. and Semaničová A.: *Numbers of edges in supermagic graphs*
3/2004 Skřivánková V. and Kočan M.: *From binomial to Black-Scholes model using the Liapunov version of central limit theorem*
4/2004 Jakubíková-Studenovská D.: *Retracts of monounary algebras corresponding to groupoids*
5/2004 Hajduková J.: *On coalition formation games*
6/2004 Fabrici I., Jendroľ S. and Semanišin G., ed.: *Czech – Slovak Conference GRAPHS 2004*
7/2004 Berežný Š. and Lacko V.: *The color-balanced spanning tree problem*
8/2004 Horňák M. and Kocková Z.: *On complete tripartite graphs arbitrarily decomposable into closed trails*
9/2004 van Aardt S. and Semanišin G.: *Non-intersecting detours in strong oriented graphs*
10/2004 Ohriska J. and Žulová A.: *Oscillation criteria for second order non-linear differential equation*
11/2004 Kardoš F. and Jendroľ S.: *On octahedral fulleroids*

2005

- 1/2005 Ceclárová K. and Vaľová V.: *The stable multiple activities problem*
2/2005 Lihová J.: *On convexities of lattices*
3/2005 Horňák M. and Woźniak M.: *General neighbour-distinguishing index of a graph*
4/2005 Mojsej I. and Ohriska J.: *On solutions of third order nonlinear differential equations*
5/2005 Ceclárová K., Fleiner T. and Manlove D.: *The kidney exchange game*

- 6/2005 Fabrici I., Jendroľ S. and Madaras T., ed.: *Workshop Graph Embeddings and Maps on Surfaces 2005*
- 7/2005 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2005*

2006

- 1/2006 Semanišinová I. and Trenkler M.: *Discovering the magic of magic squares*
- 2/2006 Jendroľ S.: *NOTE – Rainbowness of cubic polyhedral graphs*
- 3/2006 Horňák M. and Woźniak M.: *On arbitrarily vertex decomposable trees*
- 4/2006 Cechlárová K. and Lacko V.: *The kidney exchange problem: How hard is it to find a donor ?*
- 5/2006 Horňák M. and Kocková Z.: *On planar graphs arbitrarily decomposable into closed trails*
- 6/2006 Biró P. and Cechlárová K.: *Inapproximability of the kidney exchange problem*
- 7/2006 Rudašová J. and Soták R.: *Vertex-distinguishing proper edge colourings of some regular graphs*
- 8/2006 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2006*
- 9/2006 Borbeľová V. and Cechlárová K.: *Pareto optimality in the kidney exchange game*
- 10/2006 Harminc V. and Molnár P.: *Some experiences with the diversity in word problems*
- 11/2006 Horňák M. and Zlámalová J.: *Another step towards proving a conjecture by Plummer and Toft*
- 12/2006 Hančová M.: *Natural estimation of variances in a general finite discrete spectrum linear regression model*

2007

- 1/2007 Haluška J. and Hutník O.: *On product measures in complete bornological locally convex spaces*