



P. J. ŠAFÁRIK UNIVERSITY
FACULTY OF SCIENCE
INSTITUTE OF MATHEMATICS
Jesenná 5, 041 54 Košice, Slovakia



J. Miškuf, R. Škrekovski, M. Tancer

**Backbone colorings and generalized
Mycielski's graphs**

IM Preprint, series A, No. 2/2008
March 2008

Backbone Colorings and Generalized Mycielski's Graphs*

Jozef Miškuf[†] Riste Škrekovski[‡] Martin Tancer[§]

Abstract

For a graph G and its spanning tree T the *backbone chromatic number*, $\text{BBC}(G, T)$, is defined as the minimum k such that there exists a coloring $c: V(G) \rightarrow \{1, 2, \dots, k\}$ satisfying $|c(u) - c(v)| \geq 1$ if $uv \in E(G)$ and $|c(u) - c(v)| \geq 2$ if $uv \in E(T)$.

Broersma et al. [1] asked whether there exists a constant c such that for every triangle-free graph G with an arbitrary spanning tree T the inequality $\text{BBC}(G, T) \leq \chi(G) + c$ holds. We answer this question negatively by showing the existence of triangle-free graphs R_n and their spanning trees T_n such that $\text{BBC}(R_n, T_n) = 2\chi(R_n) - 1 = 2n - 1$.

In order to answer the question we obtain a result of independent interest. We modify the well known Mycielski's construction and construct triangle-free graphs J_n , for every integer n , with chromatic number n and 2-tuple chromatic number $2n$ (here 2 can be replaced by any integer t).

Keywords: backbone coloring, graph coloring, generalized Mycielski's construction, triangle-free graph.

MSC: 05C15

*This work was partially supported by the Czech-Slovenian bilateral research project 15/2006-2007 and by the Slovak-Slovenian bilateral research project 1/2006-2007.

[†]Institute of Mathematics, Faculty of Science, University of Pavol Jozef Šafárik, Jesenná 5, 041 54 Košice, Slovakia. Supported in part by Slovak Research and Development Agency under the contract No. APVV-0007-07. E-mail: jozef.miskuf@upjs.sk

[‡]Department of Mathematics, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia. Supported in part by ARRS Research Program P1-0297.

[§]Institute for Theoretical Computer Science (supported by project 1M0545 of The Ministry of Education of the Czech Republic), Faculty of Mathematics and Physics, Charles University, Malostranské nám. 25, 118 00 Prague, Czech Republic. E-mail: tancer@kam.mff.cuni.cz

1 Introduction

1.1 Backbone colorings

The backbone coloring problem is related to frequency assignment problems in the following way, the transmitters are represented by the vertices of a graph and they are adjacent in the graph if the corresponding transmitters are close enough or transmitters are strong enough. The problem is to assign frequency channels to the transmitters in such a way that the interference is kept at an "acceptable" level. One way of putting these requirements together is following: Given graphs G_1, G_2 such that G_1 is a spanning subgraph of G_2 . Determine a coloring of G_2 that satisfies certain restriction of one type in G_1 and of the other type in G_2 .

In this concept, backbone colorings were introduced and motivated and put into a general framework of related coloring problems in [1]. Let us recall some basic definitions. In the sequel we deal with undirected simple graphs, i.e. without loops and/or multiedges. By the symbol $[n]$ we understand the set $\{1, 2, \dots, n\}$, by the symbol $\chi(G)$ the chromatic number of G , and by the symbol $G[W]$ the subgraph induced by the vertex set $W \subseteq V(G)$. For a graph G , we define a coloring $\nu : V \rightarrow \{1, 2, \dots, k\}$ to be a *backbone k -coloring* of a graph G with a backbone graph $H \subseteq G$, if for every two different vertices u and v of G , it holds

- $|\nu(u) - \nu(v)| \geq 1$, if $uv \in E(G) \setminus E(H)$;
- $|\nu(u) - \nu(v)| \geq 2$, if $uv \in E(H)$.

The minimum k for which G with backbone H admits a backbone k -coloring is called the *backbone chromatic number* of G with backbone H . It is denoted by $\text{BBC}(G, H)$. In this paper we consider only the case when a backbone graph H is acyclic.

We refer to several results concerning backbone colorings of graphs. The connection between the backbone chromatic number and the chromatic number is studied in [1]. The authors showed that the backbone chromatic number of a graph G is at most $2\chi(G) - 1$, while they provided examples where this bound is attained. To show this inequality it is sufficient to color the graph G with colors $1, 3, \dots, 2\chi(G) - 1$. The decision problem if there exists a backbone coloring of a graph G with backbone tree T with l colors is NP-complete for $l \geq 5$. Broersma et al. in [3] showed that the backbone chromatic number of planar graphs with backbone matchings is at most six. Other results on backbone colorings appear in [2, 4, 9].

We deal with the intriguing question posed by Broersma et al. [1].

Question 1.1. *Does there exist a constant c such that $\text{BBC}(G, T) \leq \chi(G) + c$ holds for every triangle-free graphs G with T being a tree?*

We will present infinite class of triangle-free graphs answering the question negatively. More precisely, for every integer n , we will show the existence of a triangle-free graph G with a backbone tree T such that $\text{BBC}(G, T) = 2\chi(G) - 1 = 2n - 1$.

1.2 Triangle-free graphs and their colorings

For integers k and t we define a t -tuple k -coloring of a graph G to be a function $c: V(G) \rightarrow \binom{[k]}{t}$ such that $c(u) \cap c(v) = \emptyset$ whenever $uv \in E(G)$. The minimum possible k that G has a t -tuple k -coloring is called the t -tuple chromatic number, denoted by $\chi_t(G)$.

The procedure of giving a negative answer to Question 1.1 will proceed in the following steps:

- Step I. For a given triangle-free graph G , we will construct an infinite triangle-free graph R_G with a backbone tree T_G such that $\text{BBC}(R_G, T_G) \geq \chi_2(G) - 1$ and $\chi(R_G) = \chi(G)$.
- Step II. For a given triangle-free graph G , we will present a Mycielski-type construction of a triangle-free graph $J(G)$ such that $\chi_2(J(G)) \geq \chi_2(G) + 2$ and $\chi(J(G)) \leq \chi(G) + 1$. In particular, it follows that $\chi(J_n) = n$ and $\chi_2(J_n) = 2n$, where $J_n = J^{n-2}(K_2)$ and K_2 is the complete graph on 2 vertices.
- Step III. From the previous two steps, $\text{BBC}(R_{J_n}, T_{J_n}) \geq 2n - 1 = 2\chi(R_{J_n}) - 1$. The graph R_{J_n} is infinite; however, by the principle of compactness there exists a finite (connected) subgraph $R_n \subseteq R_{J_n}$ such that $\text{BBC}(R_n, T_{J_n}[V(R_n)]) \geq 2n - 1$. $T_{J_n}[V(R_n)]$ is a subforest of R_n thus it can be extended to a spanning tree T_n of R_n . We know that $\text{BBC}(R_n, T_n) \geq 2n - 1$ and $\chi(R_n) \leq \chi(R_{J_n}) = n$. Actually, equalities hold since $\text{BBC}(G, T) \leq 2\chi(G) - 1$ for any graph G with backbone T .

The construction from Step I follows an idea of Broersma et al. [1]. It will be described in Section 2.

The crucial step is Step II. Its task is to construct a triangle-free graph J_n , for every integer n , such that $\chi(J_n) = n$ and $\chi_2(J_n) = 2n$. This construction may be of an independent interest. The well known *fractional chromatic number* of a graph G is defined as

$$\chi_f = \inf_{t \in \mathbb{N}} \frac{\chi_t(G)}{t}.$$

Since $\chi(G) \geq \frac{\chi_2(G)}{2} \geq \chi_f(G)$, it would be natural to look for a class, say F_n , of triangle-free graphs such that $\chi(F_n) = \chi_f(F_n) = n$. Unfortunately, we are not aware of such a class of graphs. There is a lot of known examples of triangle-free

graphs with a large chromatic number, however it seems that none of them is the case. We just give a note on some of them:

Larsen, Propp and Ullman [7] proved that a Mycielski graph with the chromatic number n has fractional chromatic number approximately equal to $\sqrt{2n}$.

Kneser graphs $KG_{3n-1,n}$ are triangle-free graphs with chromatic number $n+1$ and fractional chromatic number $\frac{3n-1}{n} < 3$. More details about colorings of Kneser graphs can be found e.g. in [6, 8].

By the probabilistic method, Erdős [5] showed that exist triangle-free graphs with an arbitrary large chromatic number actually proving that the independence number of such graphs is small (and thus even fractional chromatic number is large). However, it does not give an intuition, how far away the chromatic number and the fractional chromatic number are.

The construction of graphs J_n is a generalization of Mycielski's construction. It will be precisely described in Section 3.

For Step III we just remark that with some extra effort it is possible to work just with finite graphs and avoid any use of the principle of compactness. However, some of the proofs would be more complicated and more technical, thus we rather prefer to work with infinite graphs.

2 Relation between backbone colorings and 2-tuple colorings

In this section, for a given graph G we construct a pair of graphs (R_G, T_G) as mentioned in Step I in the introduction.

Definition 2.1. Let G be a graph. We define a pair of infinite graphs (R_G, T_G) in the following way:

1. The graph R_G is the OR-product of G and \mathbb{N} (as independent set), i.e.
 - $V(R_G) = V(G) \times \mathbb{N}$;
 - $E(R_G) = \{(v_1, n_1), (v_2, n_2)\} \mid v_1 v_2 \in E(G)\}$.
2. T_G is a spanning tree of G defined recursively in the following way: Fix a good linear ordering \preceq on $V(R_G)$. Let T_G^1 is the graph with just one vertex – the smallest element of $V(R_G)$. Suppose that T_G^i is already defined. Let us define $T_G^{i+1} \supset T_G^i$: First, we call *occupied* all the vertices of T_G^i . Gradually for every vertex $(v, k) \in T_G$ (from the smallest one to the largest one) and for every edge $uv \in E(G)$, choose j such that (u, j) is the smallest possible unoccupied vertex and add this vertex among the occupied vertices. Moreover, add the vertex (u, j) among the vertices of T_G^{i+1} and the edge $\{(v, k), (u, j)\}$ among the edges of T_G^{i+1} . Finally, we define $T_G = \bigcup_{i \in \mathbb{N}} T_G^i$.




Figure 1: The path P_3 and the pair (R_{P_3}, T_{P_3}) , where the ordering \preceq is chosen so that vertices with higher altitude in the figure are smaller in \preceq . Labels at vertices are the smallest such i that the corresponding vertex belongs to $T_{P_3}^i$.

An example of the construction is depicted on Figure 1. Notice, that the spanning tree T_G is not defined uniquely; however, for our purposes it is not important to have a unique definition. The following lemma easily follows from the construction.

Lemma 2.2. *For any graph G , the pair (R_G, T_G) has the following properties:*

1. T_G is a spanning tree of R_G .
2. If G is triangle-free then R_G is triangle-free.
3. For every vertex (v, j) of R_G and for every edge uv of G , there exists an integer j' such that $\{(v, j), (u, j')\}$ is an edge of T_G .

□

The following proposition relates the 2-tuple chromatic number of a graph G and the backbone chromatic number of the pair (R_G, T_G) .

Proposition 2.3. *Let G be a graph. Then*

1. $\chi(R_G) = \chi(G)$;
2. $\text{BBC}(R_G, T_G) \geq \chi_2(G) - 1$.

Proof. We prove each of the claims separately:

1. The graph $R_G[V(G) \times \{1\}]$ is isomorphic to G , hence $\chi(G) \leq \chi(R_G)$.

On the other hand, any coloring of G induces a coloring of R_G : For every $v \in V(G)$ and $n \in \mathbb{N}$ the vertices $(v, n) \in V(R_G)$ are assigned with the color of v . Hence $\chi(R_G) \leq \chi(G)$.

2. First, observe that $\text{BBC}(R_G, T_G)$ is finite since $\text{BBC}(R_G, T_G) \leq 2\chi(R_G) - 1 = 2\chi(G) - 1$. Let $k = \text{BBC}(R_G, T_G)$ and let ν be a backbone k -coloring of (R_G, T_G) . Our goal will be to construct a 2-tuple $(k+1)$ -coloring c of G . First, we define a function $c' : V(G) \rightarrow 2^{[k]} \setminus \{\emptyset\}$:

$$c'(v) = \{n \in [k] \mid \text{exists } j \in \mathbb{N} : \nu(v, j) = n\}.$$

Now, we define a function $c : V(G) \rightarrow \binom{[k+1]}{2}$ in the following way:

- $c(v) = \{i, i+1\}$ if $c'(v) = \{i\}$.
- $c(v)$ is any 2-element subset of $c'(v)$ if $|c'(v)| \geq 2$.

It remains to show that c is a 2-tuple coloring of G . First, observe that $c'(u) \cap c'(v) = \emptyset$ for every $uv \in E(G)$, since $\{(u, j_1), (v, j_2)\} \in E(R_G)$ for every $j_1, j_2 \in \mathbb{N}$. For any $uv \in E(G)$, we will show that $c(u) \cap c(v) = \emptyset$ by considering three cases:

$|c'(u)| \geq 2$ and $|c'(v)| \geq 2$: Since $c'(u) \cap c'(v) = \emptyset$, we infer $c(u) \cap c(v) = \emptyset$.

$c'(u) = \{i\}$ and $|c'(v)| \geq 2$ (or vice versa): Since $c'(u) \cap c'(v) = \emptyset$, we infer $i \notin c(v)$. It remains to show that $i+1 \notin c(v) \subseteq c'(v)$. For a contradiction, suppose that $i+1 \in c'(v)$. Let $(v, j) \in R_G$ be a vertex such that $\nu(v, j) = i+1$ and (u, j') be its neighbor in T_G due to Lemma 2.2(3). Then $\nu(u, j') = i$ since $c'(u) = \{i\}$. It contradicts the fact that ν is a backbone coloring of (R_G, T_G) .

$c'(u) = \{i_1\}$ and $c'(v) = \{i_2\}$: Since $c'(u) \cap c'(v) = \emptyset$, we have $i_1 \neq i_2$. Thus without loss of generality, we can assume that $i_1 < i_2$. Moreover, $i_1 + 1 \notin \{i_2\}$ from a similar reason as in the previous case. Thus, $i_1 + 1 < i_2$ implies that $c(u) \cap c(v) = \emptyset$.

□

3 Mycielski-type construction

Mycielski [10] was among the first authors who showed the existence of triangle-free graphs with arbitrarily large chromatic number. We wish, in addition, to




Figure 2: An example of Construction D. In the graph G , the subgraph $G[W]$ is indicated by a thick line; circular vertices belong to A_1 and square vertices belong to A_2 .

relate the chromatic number and the 2-tuple chromatic number. More precisely, we will show that for every $n \in \mathbb{N}$ there exists a triangle-free graph which chromatic number is n and 2-tuple chromatic is $2n$. We will present a construction that increases the chromatic number by 1, the 2-tuple chromatic number by 2, and preserves the property being triangle-free. The construction has several steps.

Construction D. Suppose that we are given a graph G ; a set $W \subseteq V(G)$ such that $G[W]$ is bipartite with parts A_1 and A_2 (possibly empty); and two vertices $z_1, z_2 \notin V(G)$.

We construct a graph¹ $D = D(G, W, z_1, z_2)$. Let² $W' = W \times \{W\}$ be a copy of W and let w' be an abbreviation for $(w, W) \in W'$ where $w \in W$. We define

$$\begin{aligned} V(D) &= V(G) \cup W' \cup \{z_1, z_2\}, \text{ and} \\ E(D) &= E(G) \\ &\cup \{w'_1 w'_2 \mid w_1, w_2 \in W, \text{ and } w_1 w_2 \in E(G)\} \\ &\cup \{v w' \mid v \in V(G) \setminus W, w \in W, \text{ and } v w \in E(G)\} \\ &\cup \{w' z_i \mid w \in A_i, i \in \{1, 2\}\}. \end{aligned}$$

An example of the construction is depicted on Figure 2. It is easy to check that the following lemma holds:

Lemma 3.1. *The graph $D = D(G, W, z_1, z_2)$ from Construction D has the following properties:*

1. *If G is triangle-free then D is triangle-free.*

¹Formally, the graph D also depends on a partition of W to A_1 and A_2 . For our purposes, it will be convenient to suppose that W is always already given with such a partition.

²For the most of the purposes $W \times \{W\}$ could be replaced by $W \times \{1\}$. However it will be convenient later on to get different copies for different W .




Figure 3: A scheme of Construction H.

2. The graph $D[(V(G) \setminus W) \cup W']$ is isomorphic to G .

□

We define another auxiliary graph:

Construction H. Suppose that we are given a graph G , and two vertices $z_1, z_2 \notin V(G)$.

We define the graph $H(G, z_1, z_2)$ in the following way: Order all $W \subseteq V(G)$ such that $G[W]$ is bipartite in a sequence W_1, W_2, \dots, W_t (and choose parts A_1, A_2 for each of them). Construct graphs $D_i = D(G, W_i, z_1, z_2)$. Finally, define

$$H = H(G, z_1, z_2) = \bigcup_{i=1}^t D_i,$$

i.e. H consists of union of sets $D(G, W_i, z_1, z_2)$ where G, z_1 and z_2 are identified in all the copies, however the sets W'_i are not identified (see Figure 3).

The following lemma is the key lemma for our construction.

Lemma 3.2. Let $H = H(G, z_1, z_2)$ be a graph from Construction H.

1. If G is triangle-free, then H is also triangle-free.
2. Let k be the 2-tuple chromatic number of G . Then, there is no 2-tuple $(k+1)$ -coloring c of H such that $c(z_1) = c(z_2)$.

Proof. The first claim easily follows from Lemma 3.1(1).

For the second claim, assume to the contrary that c is a 2-tuple $(k+1)$ -coloring of H such that $c(z_1) = c(z_2) = \{k, k+1\}$. Recall that H contains G . Let $W = \{v \in V(G) \mid c(v) \cap \{k, k+1\} \neq \emptyset\}$. It is easy to see that $G[W]$ is a bipartite, thus there exists $i \in [t]$ (where t is defined as in Construction H) such




Figure 4: A scheme of Construction J.

that $W = W_i$. The graph $G' = H[(V(G) \setminus W_i) \cup W'_i]$ is isomorphic to G according Lemma 3.1(2) (where $W'_i = W_i \times \{W_i\}$ is defined as in Construction D).

We claim that $c(v) \cap \{k, k+1\} = \emptyset$ for every $v \in V(G')$: If $v \in V(G) \setminus W_i$ then it follows from the definition of $W = W_i$, if $v \in W'_i$ then either z_1 or z_2 is a neighbor of v . Thus c restricted to G' is a 2-tuple $(k-1)$ -coloring of a graph isomorphic to G contradicting the assumptions of the lemma. \square

Finally, for a graph G we define the graph $J(G)$ that will satisfy our requirements.

Construction J. Let G be a graph, $k = \chi_2(G)$, $r = \binom{k+1}{2} + 1$, and let Z be the graph with $V(Z) = \{z_1, z_2, \dots, z_r\}$ and $E(Z) = \emptyset$. For every $i, j \in [r]$, $i \neq j$, let G_{ij} be an isomorphic copy of G , formally, $V(G_{ij}) = V(G) \times \{\{i, j\}\}$ and $E(G_{ij}) = \{(u, \{i, j\}), v, \{i, j\}\} \mid uv \in E(G)\}$. Then, we define

$$J(G) = \bigcup_{\{i, j\} \subset [r]} H(G_{ij}, z_i, z_j),$$

i.e., $J(G)$ consists of an independent set Z where between each two vertices z_i, z_j of Z there is inserted a copy of $H(G, z_i, z_j)$.

Theorem 3.3. *Let G be a graph. The graph $J(G)$ satisfies the following properties:*

1. *If G is triangle-free then $J(G)$ is also triangle-free.*
2. $\chi_2(J(G)) \geq \chi_2(G) + 2$.
3. $\chi(J(G)) \leq \chi(G) + 1$.

In fact it is not difficult to derive that $\chi_2(J(G)) = \chi_2(G) + 2$ and $\chi(J(G)) = \chi(G) + 1$, but we will not need it for our purposes.

Proof. We prove each of the claims separately:

1. This claim follows from Lemma 3.2(1) and from the fact that no two z_i and z_j are adjacent in $H(G_{ij}, z_i, z_j)$.
2. We use the notation from Construction J. Let $k = \chi_2(G)$. We will show that there is no 2-tuple $(k + 1)$ -coloring c of $J(G)$. For a contradiction, suppose that such c exists. From the pigeonhole principle, there are $i, j \in [r]$ such that $c(z_i) = c(z_j)$. But it contradicts Lemma 3.2(2) for $H = H(G_{ij}, z_i, z_j)$.
3. Again, we use the notation from Construction J. Let $l = \chi(G)$, we will show that there is an $(l + 1)$ -coloring γ of $J(G)$. First, we color all the vertices of Z with color $l + 1$, i.e. $\gamma(Z) = \{l + 1\}$. Then it is sufficient to color every $H(G_{ij}, z_i, z_j)$ separately. For notational convenience, we will color $H(G, z_1, z_2)$ following Construction H so that $\gamma(z_1) = \gamma(z_2) = l + 1$. The graph G is l -colorable, hence the coloring γ can be extended to G so that γ is a coloring of G using only colors $1, 2, \dots, l$. Finally, for $i \in [t]$ and for $(w, W_i) \in W_i \times \{W_i\}$ we define $\gamma(w, W_i) = \gamma(w)$. It is easy to check that γ is an $(l + 1)$ -coloring of $H(G, z_1, z_2)$.

□

Corollary 3.4. *For every $n \in \mathbb{N}$ there exists a (connected) triangle-free graph J_n such that $\chi(J_n) = n$ and $\chi_2(J_n) = 2n$.*

Proof. Let J_1 is the graph consisting of a single vertex. For $n \geq 2$, let $J_n = J^{n-2}(K_2)$. Theorem 3.3 implies that $\chi(J_n) \leq n$ and $\chi_2(J_n) \geq 2n$, however, it is easy to see that $\chi_2(G) \leq 2\chi(G)$ for any graph G . □

The proof of the following corollary, answering negatively Question 1.1, is explicitly written in the introduction (Step III):

Corollary 3.5. *For every $n \in \mathbb{N}$ there exists a (finite) triangle-free graph R_n and its spanning tree T_n such that $\text{BBC}(R_n, T_n) = 2\chi(R_n) - 1 = 2n - 1$.*

□

4 Conclusion

We showed the existence of triangle-free graphs R_n such that their backbone colorings (with suitable spanning tree) need $2\chi(R_n) - 1 = 2n - 1$ colors. However, these graphs contain 4-cycles. For further research, it could be interesting to describe the behavior of maximum possible backbone number for graphs with given chromatic number χ and given girth g .

The construction of a graph J_n can be for every $t \geq 2$ generalized in an obvious way to get triangle-free graphs J_n^t such that $\chi(J_n^t) = n$ and $\chi_t(J_n^t) = tn$ (compare

with Corollary 3.4). In a bit more detail, to construct graphs J_n^t , consider t -colorable subgraphs W instead of bipartite subgraphs in Constructions D and H and put $r = \binom{k+t-1}{t} + 1$ in Construction J. Motivated by these results we pose the following conjecture:

Conjecture. *For every $n \in \mathbb{N}$ there exists a finite triangle-free graph F_n such that $\chi(F_n) = \chi_f(F_n) = n$.*

References

- [1] H. Broersma, F. V. Fomin, P. A. Golovach, and G. J. Woeginger. Backbone colorings for graphs: Tree and path backbones. *J. Graph Theory*, 55(2):137–152, 2007.
- [2] H. J. Broersma, J. Fujisava, L. Marchal, D. Paulusma, A. N. M. Salman, and K. Yoshimoto. λ -backbone colorings along pairwise disjoint stars and matchings. Manuscript, 2007.
- [3] H. J. Broersma, J. Fujisawa, and K. Yoshimoto. Backbone colorings along perfect matchings. Memorandum 1706, Enschede, 2003.
- [4] H. J. Broersma, L. Marchal, and D. Paulusma. Backbone colorings along stars and matchings in split graphs: their span is close to the chromatic number. Manuscript, 2007.
- [5] P. Erdős. Graph theory and probability. *Canad. J. Math.*, 11:34–38, 1959.
- [6] P. Hell and J. Nešetřil. *Graphs and homomorphisms*. Oxford University Press, 2004.
- [7] M. Larsen, J. Propp, and D. Ullman. The fractional chromatic number of Mycielski's graphs. *J. Graph Theory*, 19:411–416, 1995.
- [8] J. Matoušek. *Using the Borsuk-Ulam Theorem*. Springer, Berlin etc., 2003.
- [9] J. Miškuf, R. Škrekovski, and M. Tancer. Backbone colorings of graphs with bounded maximum degree. Manuscript, 2008.
- [10] J. Mycielski. Sur le coloriage des graphes. *Colloq. Math.*, 3:161–162, 1955.

Recent IM Preprints, series A

2003

- 1/2003 Ceclárová K.: *Eigenvectors of interval matrices over max-plus algebra*
2/2003 Mihók P. and Semanišin G.: *On invariants of hereditary graph properties*
3/2003 Ceclárová K.: *A problem on optimal transportation*

2004

- 1/2004 Jendroľ S. and Voss H.-J.: *Light subgraphs of graphs embedded in the plane and in the projective plane – survey*
2/2004 Drajnová S., Ivančo J. and Semaničová A.: *Numbers of edges in supermagic graphs*
3/2004 Skřivánková V. and Kočan M.: *From binomial to Black-Scholes model using the Liapunov version of central limit theorem*
4/2004 Jakubíková-Studenovská D.: *Retracts of monounary algebras corresponding to groupoids*
5/2004 Hajduková J.: *On coalition formation games*
6/2004 Fabrici I., Jendroľ S. and Semanišin G., ed.: *Czech – Slovak Conference GRAPHS 2004*
7/2004 Berežný Š. and Lacko V.: *The color-balanced spanning tree problem*
8/2004 Horňák M. and Kocková Z.: *On complete tripartite graphs arbitrarily decomposable into closed trails*
9/2004 van Aardt S. and Semanišin G.: *Non-intersecting detours in strong oriented graphs*
10/2004 Ohriska J. and Žulová A.: *Oscillation criteria for second order non-linear differential equation*
11/2004 Kardoš F. and Jendroľ S.: *On octahedral fulleroids*

2005

- 1/2005 Ceclárová K. and Vaľová V.: *The stable multiple activities problem*
2/2005 Lihová J.: *On convexities of lattices*
3/2005 Horňák M. and Woźniak M.: *General neighbour-distinguishing index of a graph*
4/2005 Mojsej I. and Ohriska J.: *On solutions of third order nonlinear differential equations*
5/2005 Ceclárová K., Fleiner T. and Manlove D.: *The kidney exchange game*
6/2005 Fabrici I., Jendroľ S. and Madaras T., ed.: *Workshop Graph Embeddings and Maps on Surfaces 2005*
7/2005 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2005*

2006

- 1/2006 Semanišinová I. and Trenkler M.: *Discovering the magic of magic squares*
2/2006 Jendroľ S.: *NOTE – Rainbowness of cubic polyhedral graphs*
3/2006 Horňák M. and Woźniak M.: *On arbitrarily vertex decomposable trees*
4/2006 Ceclárová K. and Lacko V.: *The kidney exchange problem: How hard is it to find a donor ?*

- 5/2006 Horňák M. and Kocková Z.: *On planar graphs arbitrarily decomposable into closed trails*
- 6/2006 Biró P. and Ceclárová K.: *Inapproximability of the kidney exchange problem*
- 7/2006 Rudašová J. and Soták R.: *Vertex-distinguishing proper edge colourings of some regular graphs*
- 8/2006 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2006*
- 9/2006 Borbeľová V. and Ceclárová K.: *Pareto optimality in the kidney exchange game*
- 10/2006 Harminc V. and Molnár P.: *Some experiences with the diversity in word problems*
- 11/2006 Horňák M. and Zlámalová J.: *Another step towards proving a conjecture by Plummer and Toft*
- 12/2006 Hančová M.: *Natural estimation of variances in a general finite discrete spectrum linear regression model*

2007

- 1/2007 Haluška J. and Hutník O.: *On product measures in complete bornological locally convex spaces*
- 2/2007 Cichacz S. and Horňák M.: *Decomposition of bipartite graphs into closed trails*
- 3/2007 Hajduková J.: *Condorcet winner configurations in the facility location problem*
- 4/2007 Kovárová I. and Mihalčová J.: *Vplyv riešenia jednej difúznej úlohy a následný rozbor na riešenie druhej difúznej úlohy o 12-tich kockách*
- 5/2007 Kovárová I. and Mihalčová J.: *Prieskum tvorivosti v žiackych riešeniach vägne formulovanej úlohy*
- 6/2007 Haluška J. and Hutník O.: *On Dobrakov net submeasures*
- 7/2007 Jendroľ S., Miškuf J., Soták R. and Škrabuláková E.: *Rainbow faces in edge colored plane graphs*
- 8/2007 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2007*
- 9/2007 Ceclárová K.: *On coalitional resource games with shared resources*

2008

- 1/2008 Miškuf J., Škrekovski R., Tancer M.: *Backbone colorings of graphs with bounded degree*