



**P. J. ŠAFÁRIK UNIVERSITY**  
**FACULTY OF SCIENCE**  
**INSTITUTE OF MATHEMATICS**  
Jesenná 5, 040 01 Košice, Slovakia



---

**V. Skřivánková and M. Juhás**

**Characterization of standard extreme  
value distributions using records**

IM Preprint, series A, No. 11/2010  
July 2010

# Characterization of standard extreme value distributions using records

Valéria Skřivánková and Matej Juhás

## Abstract

The paper deals with characterization of standard Gumbel distribution and standard Fréchet distribution and was motivated by [4], where the Weibull distribution is characterized. We present criterions using the independence of some suitable functions of lower records in a sequence of independent identically distributed random variables  $\{X_n, n \geq 1\}$ .

**Keywords:** record value, characterization, independence, Gumbel distribution, Fréchet distribution

**Mathematics Subject Classification 2000:** Primary 60E05; Secondary 60E10, 62E10.

## 1 Introduction

Consider the sequence  $\{X_n, n \geq 1\}$  of independent, identically distributed (iid) random variables with common absolutely continuous distribution function  $F$  and probability density function (pdf)  $f$ . Random variable  $X_n$  is a lower record if  $X_n < \min\{X_1, X_2, \dots, X_{n-1}\}$ . By convention  $X_1$  is a record value. Let  $\{T_n, n \geq 1\}$  be the lower record times at which record values occur. We consider discrete time and define  $T_1 = 1$  and  $T_n = \min\{i, i > T_{n-1}, X_i < X_{T_{n-1}}, n > 1\}$ . So the sequence  $\{L_n, n \geq 1\} = \{X_{T_n}, n \geq 1\}$  is a sequence of lower record values. The joint pdf of  $L_m$  and  $L_n$  and the pdf of  $L_n$  are (see [2])

$$f_{m,n}(x, y) = \frac{(H(x))^{m-1}}{\Gamma(m)} h(x) \frac{(H(y) - H(x))^{n-m-1}}{\Gamma(n-m)} f(y), \quad (1)$$

for  $1 \leq m < n$  and  $-\infty < y < x < \infty$ ,

$$f_n(x) = \frac{(H(x))^{n-1}}{\Gamma(n)} f(x), \text{ for } n \geq 1, \quad -\infty < x < \infty, \quad (2)$$

where  $H(x) = -\ln F(x)$  and  $h(x) = -\frac{d}{dx}H(x)$  are the corresponding hazard function and hazard rate. Similar relations for upper records are proved in [5].

A continuous random variable  $X$  is distributed according to standard Gumbel law if its distribution function  $F$  is of the form

$$F(x) = \exp(-e^{-x}), \quad -\infty < x < \infty. \quad (3)$$

We say that a continuous random variable  $X$  is distributed according to standard Fréchet law if its distribution function  $F$  is of the form

$$F(x) = \begin{cases} \exp(-x^{-\alpha}), & x > 0 \text{ and } \alpha > 0, \\ 0, & \text{otherwise.} \end{cases} \quad (4)$$

Some characterizations by the independence of upper record values are known. For example Ahsanullah characterized the exponential distribution (see [2]), Chang in 2007 characterized Pareto distribution (see [3]) and Lee and Lim in 2010 Weibull distribution (see [4]).

In this paper, we will give some characterizations of the standard Gumbel and the standard Fréchet distribution by the independence of certain functions of lower record values. For proving our main theorems we need the following Lemma given by Ahsanullah ([2] p. 78).

**Lemma 1.1.** *Let  $F_1(x)$  be an absolutely continuous function and  $F_1(x) < 1$ , for all  $x > 0$ . Suppose that  $q(u, v) = H_1(u + v) - H_1(u)$  and*

$$k(u, v) = (q(u, v))^r \exp(-q(u, v)) \frac{\partial}{\partial v} q(u, v), \quad r \geq 0,$$

where  $k(u, v) \neq 0$ , and  $\frac{\partial}{\partial v} q(u, v) \neq 0$  for any positive  $u$  and  $v$ . If  $k(u, v)$  is independent of  $u$  then  $q(u, v)$  is a function of  $v$  only.

## 2 Characterization of Gumbel distribution

**Theorem 2.1.** *Let  $\{X_n, n \geq 1\}$  be a sequence of iid random variables with absolutely continuous distribution function  $F(x)$  and pdf  $f(x)$ , where  $0 < F(x) < 1$  for all  $x$  real. Then  $F(x) = \exp(-e^{-x})$ ,  $-\infty < x < \infty$  if and only if  $U = e^{-L_m}$  and  $V = e^{-L_n} - e^{-L_m}$  are independent for  $1 \leq m < n$ .*

*Proof.* Let  $F(x) = \exp(-e^{-x})$ ,  $-\infty < x < \infty$ . Then  $H(x) = e^{-x}$  and according to (1) the joint pdf of  $L_m$  and  $L_n$  is for  $-\infty < y < x < \infty$  and  $1 \leq m < n$

$$f_{m,n}(x, y) = \frac{(e^{-x})^{m-1}}{\Gamma(m)} e^{-x} \frac{(e^{-y} - e^{-x})^{n-m-1}}{\Gamma(n-m)} e^{-y} e^{-e^{-y}}. \quad (5)$$

Consider the transformation

$$\begin{aligned} t : \begin{pmatrix} L_m \\ L_n \end{pmatrix} &\rightarrow \begin{pmatrix} e^{-L_m} \\ e^{-L_n} - e^{-L_m} \end{pmatrix} = \begin{pmatrix} U \\ V \end{pmatrix} \\ \tau : \begin{pmatrix} U \\ V \end{pmatrix} &\rightarrow \begin{pmatrix} -\ln U \\ -\ln(U + V) \end{pmatrix}. \end{aligned} \tag{6}$$

The Jacobian of the transformation  $\tau$  is

$$|J_\tau| = \frac{1}{U(U + V)}, \tag{7}$$

thus for the joint pdf of transformed  $U$  and  $V$  we get

$$f_{U,V}(u, v) = \frac{u^{m-1}v^{n-m-1}}{\Gamma(m)\Gamma(n-m)} e^{-u}e^{-v}, \text{ for } u > 0, v > 0. \tag{8}$$

The marginal pdf of  $U$  is given by

$$f_U(u) = \int_0^\infty f_{U,V}(u, v)dv = \frac{u^{m-1}}{\Gamma(m)} e^{-u}, \quad u > 0, \tag{9}$$

and the marginal pdf of  $V$  is

$$f_V(v) = \int_0^\infty f_{U,V}(u, v)du = \frac{v^{n-m-1}}{\Gamma(n-m)} e^{-v}, \quad v > 0. \tag{10}$$

From (8), (9) and (10) we obtain  $f_U(u)f_V(v) = f_{U,V}(u, v)$ . Hence  $U$  and  $V$  are independent.

Now we will prove the sufficient condition. So we suppose that  $U, V$  are independent for  $1 \leq m < n$ . The joint pdf  $f_{m,n}(x, y)$  of variables  $L_m, L_n$  is given by (1). Consider the transformation (6) with Jacobian (7). Then for the joint pdf of transformed  $U, V$  we get

$$\begin{aligned} f_{U,V}(u, v) &= \frac{[H(-\ln u)]^{m-1}[H(-\ln(u+v)) - H(-\ln u)]^{n-m-1}}{\Gamma(m)\Gamma(n-m)} \times \\ &\times \frac{h(-\ln u)f(-\ln(u+v))}{u(u+v)}, u > 0, v > 0 \end{aligned}$$

and the pdf of transformed  $U$  by (2) is

$$f_U(u) = \frac{[H(-\ln u)]^{m-1}}{\Gamma(m)} \cdot \frac{f(-\ln u)}{u}, u > 0.$$

Since  $U$  and  $V$  are independent, for the pdf of  $V$  holds

$$\begin{aligned} f_V(v) &= \frac{[H(-\ln(u+v)) - H(-\ln(u))]^{n-m-1}}{\Gamma(n-m)F(-\ln u)} \cdot \frac{f(-\ln(u+v))}{u+v} \\ &= \frac{[H(-\ln(u+v)) - H(-\ln(u))]^{n-m-1}}{\Gamma(n-m)F(-\ln u)} \cdot \frac{-\partial}{\partial v} F(-\ln(u+v)), \\ &u > 0, v > 0. \end{aligned}$$

If we denote  $F_1(x) = F(-\ln x)$  and  $H_1(x) = H(-\ln x)$ , for all  $x > 0$ , then

$$f_V(v) = \frac{[H_1(u+v) - H_1(u)]^{n-m-1}}{\Gamma(n-m)} \cdot \frac{-\partial}{\partial v} \left( \frac{F_1(u+v)}{F_1(u)} \right). \quad (11)$$

Now let

$$H_1(u+v) - H_1(u) = -\ln \frac{F_1(u+v)}{F_1(u)} = q(u, v).$$

Writing (11) in terms of  $q(u, v)$  as in Lemma 1.1 we get

$$f_V(v) = \frac{1}{\Gamma(n-m)} (q(u, v))^{n-m-1} \exp(-q(u, v)) \frac{\partial}{\partial v} q(u, v). \quad (12)$$

By the independence of  $U$  and  $V$ , we have for all  $v > 0$  that  $f_V(v)$  defined by (12) is independent of  $u$ . So if we use the Lemma 1.1 we get that  $q(u, v)$  is a function of  $v$  only. Hence

$$G(v) = q(u, v) = -\ln \frac{F_1(u+v)}{F_1(u)}.$$

Now we have to solve a functional equation of the form

$$G_1(v)F_1(u) = F_1(u+v), \quad (13)$$

where  $G_1(v) = \exp(-G(v))$ . The only continuous solution of (13) is (see [1])  $F_1(x) = ae^{cx}$ ,  $x > 0$ , where  $a$  and  $c$  are arbitrary constants. Then

$$ae^{cx} = F_1(x) = F(-\ln x).$$

If we use a substitution  $x = e^{-y}$ ,  $-\infty < y < \infty$ , the boundary conditions  $F(-\infty) = 0$ ,  $F(\infty) = 1$  and choice  $c = -1$ , then we get

$$F(x) = \exp(-e^{-x}), \quad -\infty < x < \infty,$$

what is the Gumbel distribution (3). Thus the proof is complete.  $\square$

### 3 Characterization of Fréchet distribution

**Theorem 3.1.** *Let  $\{X_n, n \geq 1\}$  be a sequence of iid random variables with absolutely continuous distribution function  $F(x)$  and pdf  $f(x)$ ,  $F(x) < 1$ , for all  $x > 0$  and  $F(0) = 0$ . Then  $F(x) = \exp(-x^{-\alpha})$ ,  $x > 0$  and  $\alpha > 0$  if and only if  $U = L_m^{-\alpha}$  and  $V = L_n^{-\alpha} - L_m^{-\alpha}$  are independent for  $1 \leq m < n$ .*

*Proof.* If  $F(x) = \exp(-x^{-\alpha})$ ,  $x > 0$ , then  $H(x) = x^{-\alpha}$  and according to (1) the joint pdf of  $L_m$  and  $L_n$  is for  $1 \leq m < n$  and  $-\infty < y < x < \infty$

$$f_{m,n}(x, y) = \frac{(x^{-\alpha})^{m-1}}{\Gamma(m)} \alpha x^{-\alpha-1} \frac{(y^{-\alpha} - x^{-\alpha})^{n-m-1}}{\Gamma(n-m)} \alpha y^{-\alpha-1} e^{-y^{-\alpha}}. \quad (14)$$

Consider the transformation

$$\begin{aligned} t : \begin{pmatrix} L_m \\ L_n \end{pmatrix} &\rightarrow \begin{pmatrix} L_m^{-\alpha} \\ L_n^{-\alpha} - L_m^{-\alpha} \end{pmatrix} = \begin{pmatrix} U \\ V \end{pmatrix} \\ \tau : \begin{pmatrix} U \\ V \end{pmatrix} &\rightarrow \begin{pmatrix} U^{-\frac{1}{\alpha}} \\ (U+V)^{-\frac{1}{\alpha}} \end{pmatrix}. \end{aligned} \quad (15)$$

The Jacobian of the transformation  $\tau$  is

$$|J_\tau| = \frac{1}{\alpha^2} [U(U+V)]^{-\frac{1+\alpha}{\alpha}}, \quad (16)$$

thus the joint pdf of transformed  $U$  and  $V$  we get in form

$$f_{U,V}(u, v) = \frac{u^{m-1} v^{n-m-1}}{\Gamma(m)\Gamma(n-m)} e^{-u} e^{-v}, \quad \text{for } u > 0, v > 0. \quad (17)$$

The marginal pdf of  $U$  is given by

$$f_U(u) = \int_0^\infty f_{U,V}(u, v) dv = \frac{u^{m-1}}{\Gamma(m)} e^{-u}, \quad u > 0, \quad (18)$$

and the marginal pdf of  $V$  we can write as

$$f_V(v) = \int_0^\infty f_{U,V}(u, v) du = \frac{v^{n-m-1}}{\Gamma(n-m)} e^{-v}, \quad v > 0. \quad (19)$$

From (17), (18) and (19) we obtain  $f_U(u)f_V(v) = f_{U,V}(u, v)$ , what means that  $U$  and  $V$  are independent.

Now we will prove the sufficient condition. Suppose that  $U, V$  are independent for  $1 \leq m < n$ . The joint pdf  $f_{m,n}(x, y)$  of variables  $L_m, L_n$  is given by (1).

Consider the transformation (15) with Jacobian (16). Then for the joint pdf of transformed  $U, V$  we obtain

$$f_{U,V}(u, v) = \frac{[H(u^{-\frac{1}{\alpha}})]^{m-1} [H((u+v)^{-\frac{1}{\alpha}}) - H(u^{-\frac{1}{\alpha}})]^{n-m-1}}{\Gamma(m)\Gamma(n-m)} \times \\ \times \frac{h(u^{-\frac{1}{\alpha}})f((u+v)^{-\frac{1}{\alpha}})[u(u+v)]^{-\frac{\alpha+1}{\alpha}}}{\alpha^2}, u > 0, v > 0$$

and the marginal pdf of transformed  $U$  we get by (2) in form

$$f_U(u) = \frac{[H(u^{-\frac{1}{\alpha}})]^{m-1}}{\Gamma(m)} \cdot \frac{f(u^{-\frac{1}{\alpha}})u^{-\frac{\alpha+1}{\alpha}}}{\alpha}, u > 0.$$

Since  $U$  and  $V$  are independent so the pdf of  $V$  is given by

$$f_V(v) = \frac{[H((u+v)^{-\frac{1}{\alpha}}) - H(u^{-\frac{1}{\alpha}})]^{n-m-1}}{\Gamma(n-m)} \cdot \frac{f((u+v)^{-\frac{1}{\alpha}})(u+v)^{-\frac{\alpha+1}{\alpha}}}{\alpha F(u^{-\frac{1}{\alpha}})} \\ = \frac{[H((u+v)^{-\frac{1}{\alpha}}) - H(u^{-\frac{1}{\alpha}})]^{n-m-1}}{\Gamma(n-m)F(u^{-\frac{1}{\alpha}})} \cdot \frac{-\partial}{\partial v} F((u+v)^{-\frac{1}{\alpha}}), \\ u > 0, v > 0.$$

If we denote  $F_1(x) = F(x^{-\frac{1}{\alpha}})$  and  $H_1(x) = H(x^{-\frac{1}{\alpha}})$  for all  $x > 0$ , then the remainder part of the proof is the same as the proof of Theorem 2.1.

For the marginal pdf  $f_V(v)$  the term (11) holds. The condition of the independence of  $U$  and  $V$  leads (using terms as in Lemma 1.1) to the functional equation (13). The only continuous solution of this equation is again  $F_1(x) = ae^{cx}$ ,  $x > 0$ , where  $a$  and  $c$  are arbitrary constants. So

$$ae^{cx} = F_1(x) = F(x^{-\frac{1}{\alpha}}).$$

Now we use a substitution  $x = y^{-\alpha}$ , for  $y > 0$  and choice  $c = -1$ . Then according to the boundary conditions  $F(0) = 0$ ,  $F(\infty) = 1$  we get

$$F(x) = \exp(-x^{-\alpha}), x > 0, \alpha > 0,$$

what is the Fréchet distribution (4). This completes the proof.  $\square$

### Acknowledgements

This work was supported by Slovak grant foundation under VEGA No. 1/0131/09 and 1/0325/10.

## References

- [1] ACZEL, J. : Lectures on Functional Equations and Their Applications. Academic Press, NY, 1966.
- [2] AHSANULLAH, M. : Record values - theory and application. University Press of America, Inc., Lanham, Maryland, USA, 2005. ISBN 0-7618-2794-3.
- [3] CHANG, S. K. : Characterizations of the Pareto distribution by the independence of record values, J. Chungcheong Math. Soc. 20 (2007), No. 1 , 51-57.
- [4] LEE, M. Y. and LIM, E. H. : On characterizations of the Weibull distribution by the independent property of record value, J. Chungcheong Math. Soc. 23 (2010), No. 2 , 245-250.
- [5] SKŘIVÁNKOVÁ, V. and JUHÁS, M. : Records in Non-life Insurance, IM Preprint, series A, No. 9/2010, June 2010.

Doc.RNDr. Valéria Skřivánková, CSc.  
Institute of Mathematics  
P. J. Šafárik University of Košice  
Jesenná 5, 040 01 Košice, Slovakia  
e-mail: valeria.skrivankova@upjs.sk

Mgr. Matej Juhás  
Institute of Mathematics  
P. J. Šafárik University of Košice  
Jesenná 5, 040 01 Košice, Slovakia  
e-mail: matej.juhas@student.upjs.sk



## Recent IM Preprints, series A

### 2007

- 1/2007 Haluška J. and Hutník O.: *On product measures in complete bornological locally convex spaces*
- 2/2007 Cichacz S. and Horňák M.: *Decomposition of bipartite graphs into closed trails*
- 3/2007 Hajduková J.: *Condorcet winner configurations in the facility location problem*
- 4/2007 Kovárová I. and Mihalčová J.: *Vplyv riešenia jednej difúznej úlohy a následný rozbor na riešenie druhej difúznej úlohy o 12-tich kockách*
- 5/2007 Kovárová I. and Mihalčová J.: *Prieskum tvorivosti v žiackych riešeniach vágne formulovanej úlohy*
- 6/2007 Haluška J. and Hutník O.: *On Dobrakov net submeasures*
- 7/2007 Jendroľ S., Miškuf J., Soták R. and Škrabuláková E.: *Rainbow faces in edge colored plane graphs*
- 8/2007 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2007*
- 9/2007 Cechlárová K.: *On coalitional resource games with shared resources*

### 2008

- 1/2008 Miškuf J., Škrekovski R. and Tancer M.: *Backbone colorings of graphs with bounded degree*
- 2/2008 Miškuf J., Škrekovski R. and Tancer M.: *Backbone colorings and generalized Mycielski's graphs*
- 3/2008 Mojsej I.: *On the existence of nonoscillatory solutions of third order nonlinear differential equations*
- 4/2008 Cechlárová K. and Fleiner T.: *On the house allocation markets with duplicate houses*
- 5/2008 Hutník O.: *On Toeplitz-type operators related to wavelets*
- 6/2008 Cechlárová K.: *On the complexity of the Shapley-Scarf economy with several types of goods*
- 7/2008 Zlámalová J.: *A note on cyclic chromatic number*
- 8/2008 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2008*
- 9/2008 Czap J. and Jendroľ S.: *Colouring vertices of plane graphs under restrictions given by faces*

### 2009

- 1/2009 Zlámalová J.: *On cyclic chromatic number of plane graphs*
- 2/2009 Havet F., Jendroľ S., Soták R. and Škrabuláková E.: *Facial non-repetitive edge-colouring of plane graphs*
- 3/2009 Czap J., Jendroľ S., Kardoš F. and Miškuf J.: *Looseness of plane graphs*
- 4/2009 Hutník O.: *On vector-valued Dobrakov submeasures*
- 5/2009 Haluška J. and Hutník O.: *On domination and bornological product measures*
- 6/2009 Kolková M. and Pócssová J.: *Metóda Monte Carlo na hodine matematiky*
- 7/2009 Borbel'ová V. and Cechlárová K.: *Rotations in the stable b-matching problem*
- 8/2009 Mojsej I. and Tartal'ová A.: *On bounded nonoscillatory solutions of third-order nonlinear differential equations*

- 9/2009 Jendroľ S. and Škrabuľáková E.: *Facial non-repetitive edge-colouring of semiregular polyhedra*
- 10/2009 Krajčiová J. and Pócssová J.: *Galtonova doska na hodine matematiky, kvalitatívne určenie veľkosti pravdepodobnosti udalostí*
- 11/2009 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2009*
- 12/2009 Hudák D. and Madaras T.: *On local properties of 1-planar graphs with high minimum degree*
- 13/2009 Czap J., Jendroľ S. and Kardoš F.: *Facial parity edge colouring*
- 14/2009 Czap J., Jendroľ S. and Kardoš F.: *On the strong parity chromatic number*

## 2010

- 1/2010 Cechlárová K. and Pillárová E.: *A near equitable 2-person cake cutting algorithm*
- 2/2010 Cechlárová K. and Jelínková E.: *An efficient implementation of the equilibrium algorithm for housing markets with duplicate houses*
- 3/2010 Hutník O. and Hutníková M.: *An alternative description of Gabor spaces and Gabor-Toeplitz operators*
- 4/2010 Žežula I. and Klein D.: *Orthogonal decompositions in growth curve models*
- 5/2010 Czap J., Jendroľ S., Kardoš F. and Soták R.: *Facial parity edge colouring of plane pseudographs*
- 6/2010 Czap J., Jendroľ S. and Voigt M.: *Parity vertex colouring of plane graphs*
- 7/2010 Jakubíková-Studenovská D. and Petrejčíková M.: *Complementary quasiorder lattices of monounary algebras*
- 8/2010 Cechlárová K. and Fleiner T.: *Optimization of an SMD placement machine and flows in parametric networks*
- 9/2010 Skřivánková V. and Juhás M.: *Records in non-life insurance*
- 10/2010 Cechlárová K. and Schlotter I.: *Computing the deficiency of housing markets with duplicate houses*